



This document is intended for publication in a journal, and is made available on the understanding that extracts or references will not be published prior to publication of the original, without the consent of the author.



United Kingdom Atomic Energy Authority

RESEARCH GROUP

Preprint

POPULATION DISTRIBUTIONS IN CHARGE EXCHANGE

I. Proton Capture Cross Sections for Levels $n=1$ through $n=15$

J. R. HISKES

Culham Laboratory,
Culham, Abingdon, Berkshire

1964

CLM - P 43

© - UNITED KINGDOM ATOMIC ENERGY AUTHORITY - 1964
Enquiries about copyright and reproduction should be addressed to the
Librarian, Culham Laboratory, Culham, Abingdon, Berkshire, England.

POPULATION DISTRIBUTIONS IN CHARGE EXCHANGE

I. Proton Capture Cross Sections for
Levels $n = 1$ Through $n = 15$.

by

JOHN R. HISKES*
U.K.A.E.A. Culham Laboratory, Abingdon, Berks., England.

A B S T R A C T

The charge exchange capture cross sections are calculated for captures into all states of the hydrogen atom for levels through $n = 15$ using the Brinkman-Kramers matrix element. These expressions are sufficiently general to simulate captures from all the elements assuming only one electron is active.

* On leave of absence from Lawrence Radiation Laboratory,
Livermore, California.

U.K.A.E.A. Research Group,
Culham Laboratory,
Nr. Abingdon,
Berks.

May, 1964 (C/18 ED)

C O N T E N T S

	<u>Page</u>
1. INTRODUCTION	1
2. QUANTITATIVE DISCUSSION	3
3. ACKNOWLEDGEMENT	13
4. REFERENCES	13

1. INTRODUCTION

Electron capture by protons passing through gases is known to result in captures into all states of the atom. Early calculations^{1,2} had shown that at high proton velocities captures into s-states predominate, and had indicated a population distribution which varied inversely with the third power of the principal quantum number. Later work^{3,4} has shown that in general the population distributions will be functions of five parameters involved in the capture process : the effective charges of both the target and the incident projectile, the principal quantum number and angular momentum of the active electron, and the relative velocity of the incident projectile and target. Although expressions for electron capture cross sections through principal quantum level $n = 4$ have been available⁴, there has been little explicit discussion in the literature of the dependence of the population distribution on the above mentioned parameters.

For the most part, discussion of excited state capture has been based on the Born matrix element, and the Brinkman-Kramers (BK) approximation to the Born expressions.^{3,4,5} While these approximations are known to give poor results for absolute values of the capture cross sections, there is some evidence to suggest that the BK matrix element will prove reliable for ratios of the capture cross sections. In recent experiments with thin hydrogen targets, Riviere and Sweetman^{6,7} have found the population distributions follow closely an n^{-3} distribution for levels $n = 9$ through $n = 23$, and agree with the BK ratio for captures into the $n = 14$ level for energies above 40 keV.

An interest in the population distributions for hydrogen has arisen recently in conjunction with experiments designed to produce thermonuclear plasmas. Beams of excited energetic hydrogen atoms are

expected to play an important role in contemporary neutral injection experiments.^{8,9,10} In these injection experiments the excited levels below $n = 6$ are not expected to contribute significantly because of their relatively short radiative lifetimes,¹¹ while those levels above about $n = 15$ are generally not of interest because of their rather low electric ionization thresholds.¹² It is this group of approximately ten levels, from $n = 6$ through $n = 15$, which, at the present time, appears to be of primary interest for neutral injection.

Earlier, an impact parameter method¹³ was used to calculate the s-state capture cross section for the ten lowest levels of hydrogen in the case of a lithium or cesium neutralizer.¹⁴ The use of lithium vapor as a neutralizer had been proposed previously as a means for enhancing the population of the H(2S) state.¹⁵ An experiment by Futch and Damm¹⁶ using protons incident on lithium vapor has shown that excited state populations for levels $n = 8, 9$, and 10 exhibit an increase similar to that indicated by the theory. Since the impact parameter method is based on physical assumptions similar to those leading to the BK matrix element, their experiment suggests that for purposes of comparing the population distributions for different neutralizers the BK approximation may be useful for energies as low as 20 keV. A difficulty with the impact parameter method as applied to charge exchange is that in general it does not as readily lead to closed expressions for the cross sections as does the BK method. Only in the limit of high energies, corresponding to small impact parameters, can the integrals be easily evaluated without recourse to numerical methods.¹⁷ For this reason, the BK matrix element would appear to be more suited as a basis for a theoretical survey of neutralizers than would the impact parameter method.

In this paper is discussed the capture cross sections for all

states through the $n = 15$ level in the case of protons incident on various atomic gases. Capture cross sections are presented which are derived using the BK matrix element and for a range of effective nuclear charge Z , target principal quantum level n , and angular momentum l , sufficient to simulate all the elements in a one electron approximation. It is believed these cross sections will aid in elucidating those regions of the periodic table where significant population enhancement might occur, and will give some indication of the magnitude of the population enhancement. In previous papers dealing with excited state calculations the relevant integrals have been evaluated by means of parametric differentiation.^{4,5,18} The extension of these calculations to the very highly excited levels using this method would involve considerable labor. Here we utilize the fact that the integrals can be expressed in terms of momentum eigenfunctions which in turn are readily generated using recursion relations.

2. QUANTITATIVE DISCUSSION

In this section is considered the electron capture cross sections for protons incident on an atomic neutralizer in which it is assumed that only a single outer electron is active. The BK expression of the cross section for capture into the state n,l of the hydrogen atom from a state ν,λ of a neutralizer in which the electron is moving initially in a Coulomb potential with effective charge eZ , is given by⁴

$$\sigma(\nu\lambda \rightarrow nl) = (2\pi)^{-1} M^2 \hbar^{-4} (v_f/v_i) \int_{-1}^{+1} |m|^2 d(\cos \theta), \quad (1)$$

where

$$|m|^2 = \left| \int \phi(\nu\lambda) (e^2 Z/r) \exp i\vec{\alpha} \cdot \vec{r} d\vec{r} \right|^2 \left| \int \phi^*(nl) \exp i\vec{\beta} \cdot \vec{s} d\vec{s} \right|^2. \quad (2)$$

The angle θ is the angle between the initial and final velocities \bar{v}_i and \bar{v}_f , and M is the reduced mass. The other quantities are defined by

$$\begin{aligned}\bar{\alpha} &= \bar{k}_f + \bar{k}_i M_1 / (M_1 + m); \\ \bar{\beta} &= -\bar{k}_i - \bar{k}_f M_2 / (M_2 + m); \\ \bar{k}_i &= \bar{v}_i \hbar^{-1} (M_1 + m) M_2 / (M_1 + M_2 + m); \\ \bar{k}_f &= \bar{v}_f \hbar^{-1} (M_2 + m) M_1 / (M_1 + M_2 + m).\end{aligned}$$

The masses M_1 , M_2 , and m are the target nuclei, proton, and electron masses, respectively.

Introducing a new integration variable

$$y = \alpha^2 a_0^2 + a^2 = \beta^2 a_0^2 + b^2,$$

where $a = Z/\nu$, $b = 1/n$, and a_0 the Bohr radius, eq.(1) can be written

$$\sigma(\nu\lambda - n1) = (4\pi a_0^2 p^2 e^4)^{-1} \int_x^\infty |\mathcal{M}|^2 dy; \quad (3)$$

here the upper limit is taken to be infinite as a matter of convenience. The quantities p^2 and x are defined by

$$\begin{aligned}p^2 &= (m v_i a_0 \hbar^{-1})^2, \\ x &= p^2/4 + (a^2 + b^2)/2 + (b^2 - a^2)^2/4p^2.\end{aligned}$$

The momentum eigen functions are defined as¹⁹

$$\Phi(nlm) = (2\pi)^{-3/2} \int \phi(nlm) e^{-i\bar{k} \cdot \bar{r}} d\bar{r}. \quad (4)$$

For a Coulomb potential these functions can be written

$$\Phi(nlm) = F(n1) Y(lm), \quad (5)$$

where the $Y(lm)$'s are the spherical harmonic and the general form for the function $F(n1)$ is given by Bethe and Salpeter.¹⁹

Writing eq.(2) as

$$|\mathcal{M}|^2 = |\psi(\nu\lambda)|^2 |\chi^*(n1)|^2$$

and comparing eqs.(2), (4), and (5) we see that

$$|\chi^*(nl)|^2 = 2\pi^2(2l+1) a_0^3 F^2(nl). \quad (6)$$

Using²⁰

$$|\psi(\nu\lambda)|^2 = (4a_0^2)^{-1} e^4 y^2 |\chi(\nu\lambda)|^2, \quad (7)$$

the expression (3) for the capture cross section can be written

$$\sigma(\nu\lambda - nl) = \pi^3 a_0^2 (2l+1)(4p^2 Z^3)^{-1} \int_x^\infty y^2 F^2(\nu\lambda) F^2(nl) dy. \quad (8)$$

Expressions for the cross sections have been obtained using eq.(8) for all n,l states through the $n = 15$ level and for ν,λ states $1s \rightarrow 7s$, $2p \rightarrow 6p$, and $3d \rightarrow 5d$, inclusive. Following Bates and Dalgarno (BD),⁴ advantage is taken of the fact that considerable compactness can be achieved by expressing these cross sections as simple integrals. Writing eq.(8) in the form

$$\sigma(\nu\lambda - nl) = \pi a_0^2 p^{-2} D(\nu\lambda - nl) \int_x^\infty G(\nu\lambda - nl) dy, \quad (9)$$

the appropriate quantities are:

$$D(1s - ns) = 2^8 Z^5 n^{-5}.$$

Writing $z = y^{-1} (y - 2b^2)$, the expressions for G become:

$$G(1s - 1s) = y^{-6};$$

$$G(1s - 2s) = y^{-6}(2z)^2$$

$$G(1s - 3s) = y^{-6}(4z^2 - 1)^2;$$

$$G(1s - 4s) = y^{-6}(8z^3 - 4z)^2;$$

$$G(1s - 5s) = y^{-6}(16z^4 - 12z^2 + 1)^2;$$

$$G(1s - 6s) = y^{-6}(32z^5 - 32z^3 + 6z)^2;$$

$$G(1s - 7s) = y^{-6}(64z^6 - 80z^4 + 24z^2 - 1)^2;$$

$$G(1s - 8s) = y^{-6}(128z^7 - 192z^5 + 80z^3 - 8z)^2;$$

$$G(1s - 9s) = y^{-6}(256z^8 - 448z^6 + 240z^4 - 40z^2 + 1)^2;$$

$$G(1s - 10s) = y^{-6}(512z^9 - 1024z^7 + 672z^5 - 160z^3 + 10z)^2;$$

$$G(1s - 11s) = y^{-6}(1024z^{10} - 2304z^8 + 1792z^6 - 560z^4 + 60z^2 - 1)^2;$$

$$G(1s - 12s) = y^{-6} (2048z^{11} - 5120z^9 + 4608z^7 - 1792z^5 + 280z^3 - 12z)^2;$$

$$G(1s - 13s) = y^{-6} (4096z^{12} - 11264z^{10} + 11520z^8 - 5376z^6 + 1120z^4 - 84z^2 + 1)^2;$$

$$G(1s - 14s) = y^{-6} (8192z^{13} - 24576z^{11} + 28160z^9 - 15360z^7 + 4032z^5 - 448z^3 + 14z);$$

$$G(1s - 15s) = y^{-6} (16384z^{17} - 53248z^{12} + 67584z^{10} - 42240z^8 + 13440z^6 - 2016z^4 + 112z^2 - 1)^2;$$

$$D(1s - np) = (3) 2^{12} Z^5 n^{-7} (n^2 - 1)^{-1}.$$

$$G(1s - 2p) = y^{-8} (y - b^2);$$

$$G(1s - 3p) = y^{-8} (y - b^2) (4z)^2;$$

$$G(1s - 4p) = y^{-8} (y - b^2) (12z^2 - 2)^2;$$

$$G(1s - 5p) = y^{-8} (y - b^2) (32z^3 - 12z)^2;$$

$$G(1s - 6p) = y^{-8} (y - b^2) (80z^4 - 48z^2 + 3)^2;$$

$$G(1s - 7p) = y^{-8} (y - b^2) (192z^5 - 160z^3 + 24z)^2;$$

$$G(1s - 8p) = y^{-8} (y - b^2) (448z^6 - 480z^4 + 120z^2 - 4)^2;$$

$$G(1s - 9p) = y^{-8} (y - b^2) (1024z^7 - 1344z^5 + 480z^3 - 40z)^2;$$

$$G(1s - 10p) = y^{-8} (y - b^2) (2304z^8 - 3584z^6 + 1680z^4 - 240z^2 + 5)^2;$$

$$G(1s - 11p) = y^{-8} (y - b^2) (5120z^9 - 9216z^7 + 5376z^5 - 1120z^3 + 60z)^2;$$

$$G(1s - 12p) = y^{-8} (y - b^2) (11264z^{10} - 23040z^8 + 16128z^6 - 4480z^4 + 420z^2 - 6)^2;$$

$$G(1s - 13p) = y^{-8} (y - b^2) (24576z^{11} - 56320z^9 + 46080z^7 - 16128z^5 + 2240z^3 - 84z)^2;$$

$$G(1s - 14p) = y^{-8} (y - b^2) (53248z^{12} - 135168z^{10} + 126720z^8 - 53760z^6 + 10080z^4 - 672z^2 + 7)^2;$$

$$G(1s - 15p) = y^{-8} (y - b^2) (114688z^{13} - 319488z^{11} + 337920z^9 - 168960z^7 + 40320z^5 - 4032z^3 + 112z)^2.$$

$$D(1s - nd) = (5) 2^{18} z^5 n^{-9} (n^2-4)^{-1} (n^2-1)^{-1}.$$

$$G(1s - 3d) = y^{-10} (y-b^2)^2;$$

$$G(1s - 4d) = y^{-10} (y-b^2)^2 (6z)^2;$$

$$G(1s - 5d) = y^{-10} (y-b^2)^2 (24z^2-3)^2;$$

$$G(1s - 6d) = y^{-10} (y-b^2)^2 (80z^3-24z)^2;$$

$$G(1s - 7d) = y^{-10} (y-b^2)^2 (240z^4-120z^2+6)^2;$$

$$G(1s - 8d) = y^{-10} (y-b^2)^2 (672z^5-480z^2+60z)^2;$$

$$G(1s - 9d) = y^{-10} (y-b^2)^2 (1792z^6-1680z^4+360z^2-10)^2;$$

$$G(1s - 10d) = y^{-10} (y-b^2)^2 (4608z^7-5376z^5+1680z^3-120z)^2;$$

$$G(1s - 11d) = y^{-10} (y-b^2)^2 (11520z^8-16128z^6+6720z^4-840z^2+15)^2;$$

$$G(1s - 12d) = y^{-10} (y-b^2)^2 (28160z^9-46080z^7+24192z^5-4480z^3+210z)^2;$$

$$G(1s - 13d) = y^{-10} (y-b^2)^2 (67584z^{10}-126720z^8+80640z^6-20160z^4+1680z^2-21)^2;$$

$$G(1s - 14d) = y^{-10} (y-b^2)^2 (159744z^{11}-337920z^9+253440z^7-80640z^5+10080z^3-336z)^2;$$

$$G(1s - 15d) = y^{-10} (y-b^2)^2 (372736z^{12}-878592z^{10}+760320z^8-295680z^6+50400z^4-3024z^2+28)^2.$$

$$D(1s - nf) = (7) 2^{22} 3^2 z^5 n^{-11} (n^2-9)^{-1} (n^2-4)^{-1} (n^2-1)^{-1}$$

$$G(1s - 4f) = y^{-12} (y-b^2)^3;$$

$$G(1s - 5f) = y^{-12} (y-b^2)^3 (8z)^2;$$

$$G(1s - 6f) = y^{-12} (y-b^2)^3 (40z^2-4)^2;$$

$$G(1s - 7f) = y^{-12} (y-b^2)^3 (160z^3-40z)^2;$$

$$G(1s - 8f) = y^{-12} (y-b^2)^3 (560z^4-240z^2+10)^2;$$

$$G(1s - 9f) = y^{-12} (y-b^2)^3 (1792z^5-1120z^3+120z)^2;$$

$$G(1s - 10f) = y^{-12} (y-b^2)^3 (5376z^6-4480z^4+840z^2-20)^2;$$

$$G(1s - 11f) = y^{-12} (y-b^2)^3 (15360z^7-16128z^5+4480z^3-280z)^2;$$

$$G(1s - 12f) = y^{-12} (y-b^2)^3 (42240z^8-53760z^6+20160z^4-2240z^2+35)^2;$$

$$G(1s - 13f) = y^{-12} (y-b^2)^3 (112640z^9-168960z^7+80640z^5-13440z^3+560z)^2;$$

$$G(1s - 14f) = y^{-12} (y-b^2)^3 (292864z^{10} - 506880z^8 + 295680z^6 - 67200z^4 + 5040z^2 - 56)^2;$$

$$G(1s - 15f) = y^{-12} (y-b^2)^3 (745472z^{11} - 1464320z^9 + 1013760z^7 - 295680z^5 + 33600z^3 - 1008z)^2.$$

$$D(1s - ng) = 2^{30} 3^4 Z^5 n^{-12} [(n-5)!] [(n+4)!]^{-1}.$$

$$G(1s - 5g) = y^{-14} (y-b^2)^4;$$

$$G(1s - 6g) = y^{-14} (y-b^2)^4 (10z)^2;$$

$$G(1s - 7g) = y^{-14} (y-b^2)^4 (60z^2 - 5)^2;$$

$$G(1s - 8g) = y^{-14} (y-b^2)^4 (280z^3 - 60z)^2;$$

$$G(1s - 9g) = y^{-14} (y-b^2)^4 (1120z^4 - 420z^2 + 15)^2;$$

$$G(1s - 10g) = y^{-14} (y-b^2)^4 (4032z^5 - 2240z^3 + 210z)^2;$$

$$G(1s - 11g) = y^{-14} (y-b^2)^4 (13440z^6 - 10080z^4 + 1680z^2 - 35)^2;$$

$$G(1s - 12g) = y^{-14} (y-b^2)^4 (42240z^7 - 40320z^5 + 10080z^3 - 560z)^2;$$

$$G(1s - 13g) = y^{-14} (y-b^2)^4 (126720z^8 - 147840z^6 + 50400z^4 - 5040z^2 + 70)^2;$$

$$G(1s - 14g) = y^{-14} (y-b^2)^4 (366080z^9 - 506880z^7 + 221760z^5 - 33600z^3 + 1260z)^2;$$

$$G(1s - 15g) = y^{-14} (y-b^2)^4 (1025024z^{10} - 1647360z^8 + 887040z^6 - 184800z^4 + 12600z^2 - 126)^2.$$

$$D(1s - nh) = (11) 2^{34} 3^2 5^2 Z^5 n^{-14} [(n-6)!] [(n+5)!]^{-1}.$$

$$G(1s - 6h) = y^{-16} (y-b^2)^5;$$

$$G(1s - 7h) = y^{-16} (y-b^2)^5 (12z)^2;$$

$$G(1s - 8h) = y^{-16} (y-b^2)^5 (84z^2 - 6)^2;$$

$$G(1s - 9h) = y^{-16} (y-b^2)^5 (448z^3 - 84z)^2;$$

$$G(1s - 10h) = y^{-16} (y-b^2)^5 (2016z^4 - 672z^2 + 21)^2;$$

$$G(1s - 11h) = y^{-16} (y-b^2)^5 (8064z^5 - 4032z^3 + 336z)^2;$$

$$G(1s - 12h) = y^{-16} (y-b^2)^5 (29568z^6 - 20160z^4 + 3024z^2 - 56)^2;$$

$$G(1s - 13h) = y^{-16} (y-b^2)^5 (101376z^7 - 88704z^5 + 20160z^3 - 1008z)^2;$$

$$G(1s - 14h) = y^{-16} (y-b^2)^5 (329472z^8 - 354816z^6 + 110880z^4 - 10080z^2 + 126)^2;$$

$$G(1s - 15h) = y^{-16} (y-b^2)^5 (1025024z^9 - 1317888z^7 + 532224z^5 - 73920z^3 + 2520z)^2.$$

$$D(1s - ni) = (13) 2^{40} 3^4 5^2 Z^5 n^{-16} [(n-7)!] [(n+6)!]^{-1}.$$

$$G(1s - 7i) = y^{-18} (y-b^2)^6;$$

$$G(1s - 8i) = y^{-18} (y-b^2)^6 (14z)^2;$$

$$G(1s - 9i) = y^{-18} (y-b^2)^6 (112z^2 - 7)^2;$$

$$G(1s - 10i) = y^{-18} (y-b^2)^6 (672z^3 - 112z)^2;$$

$$G(1s - 11i) = y^{-18} (y-b^2)^6 (3360z^4 - 1008z^2 + 28)^2;$$

$$G(1s - 12i) = y^{-18} (y-b^2)^6 (14784z^5 - 6720z^3 + 504z)^2;$$

$$G(1s - 13i) = y^{-18} (y-b^2)^6 (59136z^6 - 36960z^4 + 5040z^2 - 84)^2;$$

$$G(1s - 14i) = y^{-18} (y-b^2)^6 (219648z^7 - 177408z^5 + 36960z^3 - 1680z)^2;$$

$$G(1s - 15i) = y^{-18} (y-b^2)^6 (768768z^8 - 768768z^6 + 221760z^4 - 18480z^2 + 210)^2;$$

$$D(1s - nk) = (15) (7!)^2 2^{36} Z^5 n^{-18} [(n-8)!] [(n+7)!]^{-1}.$$

$$G(1s - 8k) = y^{-20} (y-b^2)^7;$$

$$G(1s - 9k) = y^{-20} (y-b^2)^7 (16z)^2;$$

$$G(1s - 10k) = y^{-20} (y-b^2)^7 (144z^2 - 8)^2;$$

$$G(1s - 11k) = y^{-20} (y-b^2)^7 (960z^3 - 144z)^2;$$

$$G(1s - 12k) = y^{-20} (y-b^2)^7 (5280z^4 - 1440z^2 + 36)^2;$$

$$G(1s - 13k) = y^{-20} (y-b^2)^7 (25344z^5 - 10560z^3 + 720z)^2;$$

$$G(1s - 14k) = y^{-20} (y-b^2)^7 (109824z^6 - 63360z^4 + 7920z^2 - 120)^2;$$

$$G(1s - 15k) = y^{-20} (y-b^2)^7 (439296z^7 - 329472z^5 + 63360z^3 - 2640z)^2.$$

$$D(1s - n1) = (17) (8!)^2 2^{40} Z^5 n^{-20} [(n-9)!] [(n+8)!]^{-1}.$$

$$G(1s - 91) = y^{-22} (y-b^2)^8;$$

$$G(1s - 101) = y^{-22} (y-b^2)^8 (18z)^2;$$

$$G(1s - 111) = y^{-22} (y-b^2)^8 (180z^2 - 9)^2;$$

$$G(1s - 121) = y^{-22} (y-b^2)^8 (1320z^3 - 180z)^2;$$

$$G(1s -131) = y^{-22} (y-b^2)^8 (7920z^4 -1980z^2 +45)^2;$$

$$G(1s -141) = y^{-22} (y-b^2)^8 (41184z^5 -15840z^3 +990z)^2;$$

$$G(1s -151) = y^{-22} (y-b^2)^8 (192192z^6 -102960z^4 +11880z^2 -165)^2.$$

$$D(1s - nm) = (19) (9!)^2 2^{44} z^5 n^{-22} [(n-10)!] [(n+9)!]^{-1}.$$

$$G(1s -n10) = y^{-24} (y-b^2)^9;$$

$$G(1s -n11) = y^{-24} (y-b^2)^9 (20z)^2;$$

$$G(1s -n12) = y^{-24} (y-b^2)^9 (220z^2 -10)^2;$$

$$G(1s -n13) = y^{-24} (y-b^2)^9 (1760z^3 -220z)^2;$$

$$G(1s -n14) = y^{-24} (y-b^2)^9 (11440z^4 -2640z^2 +55)^2;$$

$$G(1s -n15) = y^{-24} (y-b^2)^9 (64064z^5 -22880z^3 +1320z)^2.$$

$$D(1s -nn) = (21) (10!)^2 2^{48} z^5 n^{-24} [(n-11)!] [(n+10)!]^{-1}.$$

$$G(1s -11n) = y^{-26} (y-b^2)^{10};$$

$$G(1s -12n) = y^{-26} (y-b^2)^{10} (22z)^2;$$

$$G(1s -13n) = y^{-26} (y-b^2)^{10} (264z^2 -11)^2;$$

$$G(1s -14n) = y^{-26} (y-b^2)^{10} (2288z^3 -264z)^2;$$

$$G(1s -15n) = y^{-26} (y-b^2)^{10} (16016z^4 -3432z^2 +66)^2.$$

$$D(1s - no) = (23) (11!)^2 2^{52} z^5 n^{-26} [(n-12)!] [(n+11)!]^{-1}.$$

$$G(1s -12o) = y^{-28} (y-b^2)^{11};$$

$$G(1s -13o) = y^{-28} (y-b^2)^{11} (24z)^2;$$

$$G(1s -14o) = y^{-28} (y-b^2)^{11} (312z^2 -12)^2;$$

$$G(1s -15o) = y^{-28} (y-b^2)^{11} (2912z^3 -312z)^2.$$

$$D(1s - nq) = (12!)^2 2^{56} 5^2 z^5 n^{-28} [(n-13)!] [(n+12)!]^{-1}.$$

$$G(1s -13q) = y^{-30} (y-b^2)^{12};$$

$$G(1s -14q) = y^{-30} (y-b^2)^{12} (26z)^2;$$

$$G(1s - 15q) = y^{-30} (y-b^2)^{12} (364z^2 - 13)^2.$$

$$D(1s - nr) = (13!)^2 2^{60} 3^3 Z^5 n^{-30} [(n-14)!] [(n+13)!]^{-1}.$$

$$G(1s - 14r) = y^{-32} (y-b^2)^{13};$$

$$G(1s - 15r) = y^{-32} (y-b^2)^{13} (28z)^2.$$

$$D(1s - nt) = (29) (14!)^2 2^{64} Z^5 n^{-32} [(n-15)!] [(n+14)!]^{-1}.$$

$$G(1s - 15t) = y^{-34} (y-b^2)^{14}.$$

Defining $w = y^{-1} (y-2a^2)$, the capture cross section for those neutralizers in which the active electron is not initially in the 1s state can be derived from:

$$D(2s - nl) = 2^{-3} D(1s - nl);$$

$$G(2s - nl) = y^{-2} (y-2a^2)^2 G(1s - nl).$$

$$D(3s - nl) = 3^{-5} D(1s - nl);$$

$$G(3s - nl) = (4w^2 - 1)^2 G(1s - nl).$$

$$D(4s - nl) = 4^{-5} D(1s - nl);$$

$$G(4s - nl) = (8w^3 - 4w)^2 G(1s - nl).$$

$$D(5s - nl) = 5^{-5} D(1s - nl);$$

$$G(5s - nl) = (16w^4 - 12w^2 + 1)^2 G(1s - nl).$$

$$D(6s - nl) = 6^{-5} D(1s - nl);$$

$$G(6s - nl) = (32w^5 - 32w^3 + 6w)^2 G(1s - nl).$$

$$D(7s - nl) = 7^{-5} D(1s - nl);$$

$$G(7s - nl) = (64w^6 - 80w^4 + 24w^2 - 1)^2 G(1s - nl).$$

$$D(2p - nl) = 2^{-3} 3^{-1} Z^2 D(1s - nl);$$

$$G(2p - nl) = y^{-2} (y-a^2) G(1s - nl).$$

$$D(3p - nl) = (2) 3^{-7} Z^2 D(1s - nl);$$

$$G(3p - nl) = y^{-2} (y-a^2) (4w)^2 G(1s - nl).$$

$$D(4p - nl) = 2^{-10} 3^{-1} 5^{-1} Z^2 D(1s - nl);$$

$$G(4p - nl) = y^{-2} (y-a^2) (12w^2 - 2)^2 G(1s - nl).$$

$$\begin{aligned}
D(5p - nl) &= (6) 5^{-7} Z^2 D(1s - nl); \\
G(5p - nl) &= y^{-2} (y-a^2) (32w^3 - 12w)^2 G(1s - nl). \\
D(6p - nl) &= 2^4 5^{-1} 6^{-7} 7^{-1} Z^2 D(1s - nl); \\
G(6p - nl) &= y^{-2} (y-a^2) (80w^4 - 48w^2 + 3)^2 G(1s - nl).
\end{aligned}$$

$$\begin{aligned}
D(3d - nl) &= 2^7 3^{-9} 5^{-1} Z^4 D(1s - nl); \\
G(3d - nl) &= y^{-4} (y-a^2)^2 G(1s - nl). \\
D(4d - nl) &= 2^{-10} 3^{-2} 5^{-1} Z^4 D(1s - nl); \\
G(4d - nl) &= y^{-4} (y-a^2)^2 (6w)^2 G(1s - nl). \\
D(5d - nl) &= 2^7 3^{-2} 5^{-9} 7^{-1} Z^4 D(1s - nl); \\
G(5d - nl) &= y^{-4} (y-a^2)^2 (24w^2 - 3)^2 G(1s - nl).
\end{aligned}$$

In labelling these states the spectroscopic notation given in Condon and Shortley⁽²¹⁾ has been used. It is easily verified that the above expressions reduce to those given by BD for states through $n = 4$.

In a subsequent paper explicit results of the population distributions for a variety of neutralizers will be presented. Here we restrict the discussion to a few semiquantitative remarks based on the general form of the capture cross sections as given by Eq.(8). Firstly, for a sufficiently high incident proton velocity and for all neutralizers we have the well-known result that the s-state capture distribution varies as n^{-3} , as deduced from the coefficient of the asymptotic form for $F^2(nl)$. Secondly, captures into the very high angular momentum states are in general not expected to contribute appreciably to the total cross section since the magnitude of $F^2(nl)$ is dominated by the coefficient

$$2^{4l} (l!)^2 [(n-l-1)!]^2 [(n+1)!]^{-2}.$$

In this connection, we note that the large resonance reported by Butler and Johnston⁽²²⁾ at $p^2 = 1$ is not reproduced in these calculations.

3. ACKNOWLEDGEMENT

The author would like to thank Dr J.B. Adams for the hospitality extended by the Culham Laboratory during the course of this visit.

4. REFERENCES

1. OPPENHEIMER, J.R., Phys. Rev. 31, 349 (1928).
2. BRINKMAN, H.C. and KRAMERS, H.A., Proc. Acad. Sci. Amsterdam, 33, 973 (1930).
3. JACKSON, J.D. and SCHIFF, H., Phys. Rev. 89, 359 (1953).
4. BATES, D.R. and DALGARNO, A. Proc. Phys. Soc. A66, 972 (1953).
5. MAPLETON, R.A. Phys. Rev. 122, 528 (1961); Phys. Rev. 126, 1477 (1962).
6. RIVIERE, A.C. and SWEETMAN, D.R. Proc. of the VI Int. Conf. on Ionization Phenomena in Gases, Paris, 1963 (Bureau des Editions Centre d'Etudes Nucleaires de Saclay) p.105.
7. RIVIERE, A.C. and SWEETMAN, D.R.. Proc. of Third Int. Conf. on Atomic Collision Processes, London, 1963.
8. SWEETMAN, D.R. Nuclear Fusion 1962 Supplement I, 279 (1962).
9. KARPUSHKINA, E.I. and GRECHUKHIN, D.L. J. Nuclear Energy, Pt C, 6, (1964).
10. HISKES, J.R. Lawrence Radiation Laboratory Report UCRL-6372, June 1961 (unpublished); Nuclear Fusion 2, 38 (1962); Phys. Rev. Letters 10, 102 (1963); Proc. of the VI Int. Conf. on Ionization Phenomena in Gases, Paris, 1963 (Bureau des Editions Centre d'Etudes Nucleaires de Saclay) p.99.
11. HISKES, J.R., TARTER, C.B. and MOODY, D.A. Phys. Rev. 133, A424 (1964).
12. BAILEY, D.S., HISKES, J.R. and RIVIERE, A.C. To be published.
13. MITTLEMAN, M.H. Phys. Rev. 122, 499 (1961).
14. HISKES, J.R. and MITTLEMAN, M.H. Controlled Thermonuclear Research Semiannual Progress Report UCRL-9969, Jan. 1962, p.128.
15. OWEN, G. (Johns Hopkins University) Private Communication. See also R. FLEISCHMANN, Nucl. Instr. and Methods, 11, 112, (1961).
16. FUTCH, A.H. and DAMM, C.C. Nuclear Fusion, 3, 124 (1963).

REFERENCES

(continued)

17. MITTLEMAN, M.H. Proc. Phys. Soc. 81, 633 (1963).
18. CORINALDESE, E. and TRAINOR, L. Nuovo Cimento, 9, 940 (1952).
19. BETHE, H.A. and SALPETER, E.E. Quantum Mechanics of One and Two Electron Atoms, (Springer-Verlag, Berlin, 1957), p.39.
20. SNEDDON, I.N. Fourier Transforms (McGraw-Hill, New York, 1951) p.380.
21. CONDON, E.U. and SHORTLEY, G.H. Theory of Atomic Spectra (Cambridge University Press, 1959) p.113.
22. BUTLER, S.T. and JOHNSTON, I.D.S. Phys. Letters, 9, 141 (1964).

