

## LH Power Losses In Front of the JET Launcher

P. Jacquet, G. Arnoux, L. Colas, L. Delpech, A. Ekedahl et al.

Citation: *AIP Conf. Proc.* **1187**, 399 (2009); doi: 10.1063/1.3273776

View online: <http://dx.doi.org/10.1063/1.3273776>

View Table of Contents: <http://proceedings.aip.org/dbt/dbt.jsp?KEY=APCPCS&Volume=1187&Issue=1>

Published by the [American Institute of Physics](#).

---

### Related Articles

A wireless handheld probe with spectrally constrained evolution strategies for diffuse optical imaging of tissue  
*Rev. Sci. Instrum.* **83**, 033108 (2012)

In vivo and simultaneous multimodal imaging: Integrated multiplex coherent anti-Stokes Raman scattering and two-photon microscopy  
*Appl. Phys. Lett.* **97**, 223702 (2010)

Suitability of using far-infrared imaging system for noncontact evaluation on working state of implantable medical devices  
*J. Appl. Phys.* **105**, 064701 (2009)

Real-time three-dimensional optoacoustic imaging using an acoustic lens system  
*Appl. Phys. Lett.* **85**, 846 (2004)

Noninvasive, low-noise, fast imaging of blood volume and deoxygenation changes in muscles using light-emitting diode continuous-wave imager  
*Rev. Sci. Instrum.* **73**, 3065 (2002)

---

### Additional information on AIP Conf. Proc.

Journal Homepage: <http://proceedings.aip.org/>

Journal Information: [http://proceedings.aip.org/about/about\\_the\\_proceedings](http://proceedings.aip.org/about/about_the_proceedings)

Top downloads: [http://proceedings.aip.org/dbt/most\\_downloaded.jsp?KEY=APCPCS](http://proceedings.aip.org/dbt/most_downloaded.jsp?KEY=APCPCS)

Information for Authors: [http://proceedings.aip.org/authors/information\\_for\\_authors](http://proceedings.aip.org/authors/information_for_authors)

### ADVERTISEMENT



AIP Advances

*Submit Now*

Explore AIP's new  
open-access journal

- Article-level metrics now available
- Join the conversation! Rate & comment on articles

# LH Power Losses In Front of the JET Launcher

P. Jacquet,<sup>a</sup> G. Arnoux,<sup>a</sup> L. Colas,<sup>b</sup> L. Delpech,<sup>b</sup> A. Ekedahl,<sup>b</sup> D. Frigione,<sup>c</sup>  
M. Goniche,<sup>b</sup> K. Kirov,<sup>a</sup> F. Leguen,<sup>d</sup> J. Mailloux,<sup>a</sup> M.-L. Mayoral,<sup>a</sup> J.  
Ongena,<sup>e</sup> V. Petržilka,<sup>f</sup> C. Portafaix,<sup>b</sup> F. Rimini,<sup>d</sup> and JET-EFDA  
contributors

*JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK.*

*<sup>a</sup>EURATOM/UKAEA Fusion Association, Culham Science Center, Abingdon, OX14 3DB, UK.*

*<sup>b</sup>Association EURATOM/CEA, IRFM, F-13108 Saint-Paul-lez Durances, France.*

*<sup>c</sup>Association EURATOM/ENEA, Frascati, Italy.*

*<sup>d</sup>EFDA Close Support Unit, Culham Science Centre, OX14 3DB, Abingdon, UK.*

*<sup>e</sup>Association EURATOM/Belgium State, LPP-ERM/KMS, B-1000 Brussels, Belgium.*

*<sup>f</sup>EURATOM/MPP.CR, 182 21 Prague 8, Czech Republic.*

**Abstract.** In recent JET experiments, Lower Hybrid (LH) power losses in the Scrape-Off Layer (SOL) were characterized using infra-red (IR) thermography. Hot spots were observed on objects intercepting the field lines passing in front of the LH launcher, i.e. on poloidal limiters and on dumplates located at the top of the tokamak; their locations being in good agreement with magnetic field line tracing using the EFIT equilibrium code. The dumplate temperature was monitored while scanning the launcher position so that the radial distance between field lines intercepting the hot spots and the launcher was increased up to 3.5 cm. The dissipation layer in front of the launcher was estimated to be at least 3.5 cm wide, in agreement with recent measurements on Tore-Supra, but not with simple models that predict a dissipation layer in the mm range.

**Keywords:** Tokamak, JET, Lower Hybrid waves, Antenna.


**PACS:** 52.55.Fa, 52.40.Fd, 52.35.Hr

## INTRODUCTION


Most of the experimental observations of localised hot spots precisely located on tokamak components magnetically connected to the launcher [1] (with in some cases connection length of few tens of m) can be explained by Electron Landau Damping (ELD) of the waves with high  $N_{\parallel}$  components ( $|N_{\parallel}| > 15$ ) directly in front of the grill, generating fast electrons in co and counter current directions with energies up to few keV [2]. However, simple wave propagation models taking into account ELD predict that these high  $|N_{\parallel}|$  waves are being absorbed within few mm [3], while recent measurements on Tore-Supra (TS) have provided evidence of fast electrons 40-50 mm in front of the grill [4]. A good knowledge of the radial extent of the LW wave dissipation in the SOL is key for the understanding of this phenomena and to estimate the amount of LH power lost through this channel. This paper describes JET

---

\* See the Appendix of F. Romanelli et al., Fusion Energy Conference 2008 (Proc. 22nd Int. FEC Geneva, 2008) LAEA, (2008)



**FIGURE 1.** JPN69582  $t=11.3$  s, IR frame showing locations where LH hot spots are observed. Also shown is the location of the LH launcher and of ICRF antennae A, B and ILA. In this example LH hot spots can be seen on 2, 3, 4, 5 and 6.




**FIGURE 2.** Cross section of plasma, from EFIT equilibrium code, JPN69582  $t=11.3$  s.


experiments in which the radial extent of the LH wave dissipation layer in front of the grill was measured. The JET LH launcher [5] is made of 12 rows of 32 reduced rectangular waveguides, fed by 6 modules of Klystrons (A-F). Depending of the plasma configuration, LH grill position, and which part of the launcher is used, LH hot spots can be observed at (see Fig. 1): the septum of ICRF antenna B and A respectively (locations 1 and 4 respectively), the protection limiters of the ICRF Iter-Like Antenna (ILA, 2 and 3), the poloidal limiter (5), the dumplates located on top of the tokamak (6). The inner divertor aprons (7). The outer divertor aprons (8). The hot spots on locations 1 to 7 correspond to electrons accelerated in the co-current direction, while location 8 intercepts electrons accelerated in the counter-current direction. Hot spots at locations 1-5 are generally observed if the bottom rows of the grill are powered. This can be understood from Fig. 2. Fast electrons travelling in the co-current direction, and generated a few centimetres in front of the bottom rows of the grill, get closer to the poloidal protections and eventually hit them when travelling from  $Z < 0$  to  $Z > 0$ . Hot spots are observed on the dumplates (6), only for plasma configurations with a pseudo X point on top of the machine. In these cases hot-spots 7 are generally not seen, and vice-versa.

## WIDTH OF THE DISSIPATION LAYER

A series of JET pulses were performed to measure the radial extent of the LH wave dissipation layer in front of the grill. The plasma parameters were the following: L-mode plasma;  $B_T=2.95$  T;  $I_P=1.75$  MA;  $q_{95}=5$ ; the separatrix-grill distance was




**FIGURE 3.** JPN77393  $t=8$  s, magnetic field lines passing in front of the grill, tracing started on a 1 cm grid, from the main poloidal limiter toward the plasma, in front of the top and the bottom rows of the grill. ‘0’ refers to lines started directly in front of the limiter, ‘3’ refers to lines started 3 cm toward the plasma.




**FIGURE 4.** JPN77393  $t=8$  s, IR view showing LH hot spots on dumplates, generated from LH grill top rows (A&B) and bottom rows (E). Encircled with dotted lines are the points used for temperature analysis in Fig. 5 and Fig. 6

approximately 9 cm, and was changed in steps of 1 cm (see Fig. 5). Gas injection close to the grill was used ( $6 \times 10^{21}$  el/s) to improve antenna coupling. The LH hot spots could be observed on the dumplates, and real time plasma shape control (XSC) was used to ensure uniform magnetic connection between the launcher and the dumplates throughout the LH phase. The connection length between the grill and the hot spots was  $\sim 20$  m. We now discuss in more details pulse No 77393, for which 1.95 MW of LH power was launched from the top rows of the grill (modules A&B) and 1.05 MW from the bottom rows (module E). The LH power density at the grill was  $\sim 24$  MW/m<sup>2</sup>. On Fig. 3, magnetic field lines are traced for this pulse at  $t=8$  s. Power dissipated in front of each row of the grill is expected to generate hot-spots on a line (each singular hot spot corresponding to protuberating surfaces), the electrons accelerated further away from the grill being intercepted further away toroidally by the dumplates, as they are closer to the pseudo X point. Also, the hot spot lines from the top and bottom rows are two separate lines with the hot spots from the bottom row being further away toroidally. This agrees well with the IR camera view on Fig. 4. Also seen on Fig. 4 is the reference ‘0’ of the bottom row (power dissipated directly in front of the grill, corresponding to point ‘0’ of bottom row hot spot line on Fig. 3). We can therefore conclude (see Fig. 3) that during the 6-8 s phase of the pulse, the top row hot spots (encircled in Fig. 4) corresponds to LH power dissipated 1.5 cm in front of the grill (which also explains their inward position relative to the bottom row hot spots). From this point, we have retracted the grill in 1 cm steps, as shown on Fig. 5. Fitting the top row hot spots temperature evolution with a semi infinite model (for graphite at 200 °C,  $k=9000$  Wm<sup>-2</sup>K<sup>-1</sup>s<sup>1/2</sup>), the heat flux on the dumplates was estimated as a function of the radial distance between magnetic field lines intercepting the analyzed hot-spots and the grill (Fig. 6). The heat flux only decreases by 20% when the grill is retracted by 2 cm, corresponding to a ‘probing’ position located 3.5 cm away from the top rows of the grill.



**FIGURE 5.** JPN77393. (a) Temperature evolution,  $Q_{\perp}$ , as a function of radial distance between field top row hot spots on dumplates (averaged over 7 lines intercepting the analyzed top row hot-spots and the grill.



**FIGURE 6.** Estimated heat flux on dumplates,  $Q_{\perp}$ , as a function of radial distance between field lines intercepting the analyzed top row hot-spots and the grill.

## CONCLUSIONS

The width of the LH power dissipation layer in front of the grill was evaluated on JET, for  $\rho_{LH} = 24 \text{ MW/m}^2$  close to the ITER target, and it was found to be at least 3.5 cm. This result, and the recent TS observations [4], are a puzzle for theory [6,7], as linear Landau damping of LH waves in front of the grill predicts interaction layer in the mm range. The maximum heat flux along the field lines was estimated to be  $\sim 10 \text{ MW/m}^2$ , this estimate will be described in more details in an extended publication.

## ACKNOWLEDGEMENTS

*The support from the JET LH system group is acknowledged. Discussions with J. Gunn and V. Fuchs were very much appreciated. This work was carried out within the European Fusion Development Agreement and was partly funded by the United Kingdom Engineering and Physical Sciences Research Council and by the European Communities under the contract of Association between EURATOM and UKAEA. One of the authors (V.P.) acknowledges partial support of the Czech Project GACR 202/07/0044. The views and opinions expressed herein do not necessarily reflect those of the European Commission.*

## REFERENCES

1. Goniche M. et al., Nucl. Fusion **38** (6), p. 919 (1998) - Rantamäki K. et al., Plasma Phys. Contr. Fusion **47** pp1101-1108 (2005).
2. Fuchs V. et al., Physics of Plasmas **11** p. 4023 (1996).
3. Rantamäki K. et al., Nucl. Fusion **40** p.1477 (2000).
4. Gunn J. et al., J. Nucl. Mat. **390-391**, p.904 (2009).
5. J.A. Dobbing et al., Proc. 17th Symp. Fusion Technology, Rome, Italy, Vol.1 p. 472, Elsevier Science, the Netherlands (1992).
6. Fuchs V. et al., 'Landau damping of the lower hybrid grill spectrum by tokamak edge electrons', these proceedings.
7. Petrzilka V. et al., 'Fast electrons generation by LH waves scattered on ponderomotive density modulations in front of LH grill', poster contribution at 2009 EPS conference.