

PAPER

## Evaluation of abrupt energy transfer among turbulent plasma structures using singular value decomposition

To cite this article: M Sasaki *et al* 2021 *Plasma Phys. Control. Fusion* **63** 025004

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Evaluation of abrupt energy transfer among turbulent plasma structures using singular value decomposition

M Sasaki<sup>1,2,3</sup> , T Kobayashi<sup>4</sup> , R O Dendy<sup>2,3,5</sup>, Y Kawachi<sup>6</sup> , H Arakawa<sup>7</sup>   
and S Inagaki<sup>1,2</sup>

<sup>1</sup> Research Institute for Applied Mechanics, Kyushu University, Kasuga 816-8580, Japan

<sup>2</sup> Research Center for Plasma Turbulence, Kyushu University, Kasuga 816-8580, Japan

<sup>3</sup> Centre for Fusion, Space and Astrophysics, Department of Physics, Warwick University, Coventry CV4 7AL, United Kingdom

<sup>4</sup> National Institute for Fusion Science, Toki 509-5292, Japan

<sup>5</sup> CCFE, Culham Science Centre, Abingdon, Oxfordshire OX14 3DB, United Kingdom

<sup>6</sup> Interdisciplinary Graduate School of Engineering Sciences, Kyushu University, Kasuga 816-8580, Japan

<sup>7</sup> Institute of Science and Engineering, Academic Assembly, Shimane University, Matsue 690-8504, Japan

E-mail: [sasaki@riam.kyushu-u.ac.jp](mailto:sasaki@riam.kyushu-u.ac.jp)

Received 20 July 2020, revised 27 October 2020

Accepted for publication 17 November 2020

Published 8 December 2020



## Abstract

A method to quantify the energy transfer among turbulent structures using singular value decomposition (SVD) is presented. We apply the method to numerical turbulence data obtained from a global plasma simulation using the Hasegawa–Wakatani fluid model, in which the Kelvin–Helmholtz instability plays a dominant role. Using the SVD method, the electrostatic potential is decomposed into a background potential deformation, a zonal flow, a coherent mode and an intermittent structure. Thus there are four key structures, as distinct from the three found in conventional theory. The kinetic energy of each structure is evaluated, and the limit cycle among them is obtained. In the limit cycle, an abrupt change of the background is found to be synchronised with the period of the zonal flow. The energy transfer function of each turbulence structure, which is defined on the basis of a vorticity equation, is evaluated. This then provides physical understanding of how the limit cycle is sustained by dynamical changes in the energy transfer among structures over its period. In addition, it is shown that the abrupt deformation of the background is caused by the non-linear self-coupling of the intermittent structure.

Keywords: structural formation, zonal flow, Kelvin–Helmholtz turbulence, singular value decomposition

(Some figures may appear in colour only in the online journal)

## 1. Introduction

Magnetically confined plasmas are non-equilibrium open systems, in which non-linear structures are observed to arise spontaneously on the largest accessible scales. These structures typically feed on microturbulence [1], and include zonal flows [2], bulk intrinsic toroidal rotation [3], and transport barriers that can arise both within the plasma [4] and at its boundary [5]. This phenomenology is of great practical

importance, because the transport of energy, momentum and particles across the plasma is conditioned by, and self-consistent with, the excitation and continuing existence of the largescale non-linear structures [6]; the latter are often referred to as secondary structures, to distinguish them from the primary microturbulence. Understanding and predicting this phenomenology, and its consequences for the local and global behaviour of the plasma, is greatly assisted if techniques applicable to low-dimensional dynamical systems [7] can be

deployed. This approach requires quantitative spatio-temporal characterisation of the salient non-linear structures and of the microturbulence, together with the couplings between them.

A pre-requisite for this approach is to have a reliable means to identify such structures, by extracting them from experimental data and from the outputs of large-scale numerical simulations. Indeed it can be a non-trivial task to determine how many distinguishable non-linear structures are present. This is important both for quantitative characterisation of the data, and to motivate or validate the adoption of low-dimensional analytical models in the predator-prey genre for turbulence phenomenology, see for example [2, 8, 9]. Notable current methods include singular value decomposition (SVD) [10, 11, 12–16] and dynamical mode decomposition (DMD) [17–19]. These methods have been widely applied in plasma physics, for example to the extraction of fluid turbulent structures [20] and the visualization of phase space structures [21]. In the DMD method, the time evolution of the field is assumed to be exponential, hence the computational cost is low. However, while the exponential assumption can work well to characterise system dynamics over relatively short timescales, beginning at any instant, it is not well adapted to system dynamics that incorporate limit cycle oscillations (LCOs) [22], which are a focus of the present paper. In plasma physics, LCOs are often observed near the threshold for local or global transitions that may also involve structure formation. Examples span electric field pulsations [23], phenomenology near the threshold for L-H transitions in global energy confinement [24–26], and edge localized modes [27]. These LCOs may significantly affect overall plasma performance, so that understanding their underlying physics is important. While the assumption of exponential time dependence in DMD treatments can be mitigated to some extent [17], DMD is not well adapted to LCOs where, as in the present paper, there is a cyclic rise and fall in the intensity of turbulence, of turbulence-driven structures, and of the coupling between them.

On the other hand, the SVD method makes no prior assumption regarding time evolution, so that the extraction of, for example, frequency chirping and phase modulation becomes possible. SVD is based on functional decomposition with respect to orthogonal basis functions which are, by construction, uncoupled to each other. This greatly assists the evaluation of the intrinsically non-linear forces and fluxes, which are expressed in relation to the decomposed structures, and which drive the fluctuations.

Here we apply the SVD method to a dataset obtained from a specific global simulation [28] of a magnetised cylindrical plasma based on the non-linear Hasegawa–Wakatani coupled two-fluid model [29] for density and electrostatic potential. In this simulation, it has been found [28] that LCOs arise spontaneously, with phenomenology dominated by the Kelvin–Helmholtz (KH) instability [30]. Using the SVD method, the electrostatic potential is decomposed into the background flow, the zonal flow, a coherent mode, and an intermittent structure. We find that the LCO between these turbulent structures is synchronised with the periodicity of the zonal flow, and involves an abrupt change in the background flow. An energy


transfer function based on the vorticity equation is introduced. This enables us to evaluate the cyclic dynamics of how the energy transfer among the turbulent structures changes during the LCO. The rest of the paper is organised as follows. Section 2 introduces the simulation dataset used in this study. In section 3, we describe how SVD is used to extract the spatio-temporal pattern of the turbulence. The dynamics of the energy transfer among the structures is evaluated in section 4. A summary is given in section 5.

## 2. Abrupt change of background and turbulence

We first introduce the dataset to which the SVD method will be applied. It relates to turbulent phenomena that arise in a global three-dimensional fluid simulation of a magnetised cylindrical plasma, based on the Hasegawa–Wakatani model [31]. The magnetic field, which is in the axial direction, is spatially uniform and constant. The turbulence and background plasma profiles are self-consistently calculated by the flux-driven simulation with a set of parameters similar to those in basic experiments in linear devices [28]. A radially inhomogeneous background flow is driven by introducing a vorticity source, such that the plasma is linearly unstable against the KH instability. The fully non-linear evolution of the KH instability is captured by the simulation. In cases with a weak vorticity source, stationary non-linear saturation is obtained. Above a certain threshold for the intensity of the vorticity source, LCO occurs [28], and we focus on this regime in the present study. Details of simulation condition and the basic characteristics of the fluctuations are given in [28].

Figure 1(a) shows a snapshot of the fluctuating component of the electrostatic potential. Noting that the electrostatic potential is intrinsically related to the  $E \times B$  bulk plasma flow, hereafter we use the terms potential or flow interchangeably, depending on the context. Here the fluctuation is translationally invariant along the axis of the cylinder; with respect to azimuthal angle, the component  $m = 1$  is dominant, where  $m$  is the azimuthal mode number. This is characteristic of the non-linear KH instability in the scenario considered [30]. The spatial pattern is dynamically modulated under the LCO, whose period is much longer than that of azimuthal rotation; further detail is given in reference [28]. Figure 1(b) shows the time evolution of the background flow, calculated from the electrostatic potential with  $m = n = 0$ ,  $V_\theta = \partial_r \phi_{0,0}$ , where  $m$  and  $n$  are the azimuthal and axial mode numbers, respectively. The radial profiles of  $V_\theta$  at different times are shown in figure 1(c). The flow curvature, which is the driving force for the KH instability, becomes strong and weak in a LCO manner. The magnitude of the velocity variation under LCO is approximately 30%.

Figure 2(a) shows the time evolution of a Fourier power spectrum with respect to azimuthal modes of the electrostatic potential. The mode spectrum for azimuthal modes up to  $m = 15$  shows that the LCO has a period  $T_{period} \sim 150$ . To further illustrate the temporal dynamics of the background potential, we construct an empirical instantaneous growth rate from the simulations, using the definition  $\gamma(t) = \partial_t \ln E_{0,0}$ , where  $E_{0,0}$  is the kinetic energy of the background flow,



**Figure 1.** (a) Snapshot of three-dimensional pattern of the electrostatic potential fluctuation at  $t = 3000$ . (b) Time evolution of the background azimuthal flow, calculated from the electrostatic potential with  $m = n = 0$ ,  $V_\theta = \partial_r \phi_{0,0}$ , where  $m$  and  $n$  are the azimuthal and axial mode numbers, respectively. (c) Radial profile of background flow at  $t = 3040, 3060, 3080$  and  $3100$ , where these times are shown in the black dashed lines in figure 1(b).

calculated from the background potential. That is,  $E_{0,0} = 2^{-1} \int |\nabla_\perp \langle \phi \rangle|^2 d^3x$ , where  $\langle \dots \rangle$  denotes the zonal average (average in the azimuthal and axial directions). The time evolution of  $\gamma(t)$  is plotted in figure 2(b) showing a sequence of rapidly rising spikes which are characteristic of this type of abrupt phenomenon [32].


### 3. Evaluation of energy transfer by SVD

Let us now apply the SVD method to the dataset described in the previous section, so as to extract the dominant spatio-temporal structures and evaluate the energy transfer among them. We shall also examine the physical origin of the abrupt increases of the background flow noted in figure 2 and identify the role of each structure in the LCO.

Because the KH turbulence is purely two-dimensional, we analyse the electrostatic potential  $\phi(r, \theta, t)$ ; here  $\phi(r, \theta, t)$  includes both the fluctuations and the background deformation. We note that the time averaged component is extracted before the analysis. Using the SVD method,  $\phi(r, \theta, t)$  is decomposed as


$$\phi(r, \theta, t) = \sum_j s_j \Psi_j(r, \theta) h_j(t). \quad (1)$$

Here, in relation to the  $j$ th mode:  $\Psi_j(r, \theta)$  describes the mode's spatial structure;  $h_j(t)$  captures its temporal evolution; and  $s_j$



**Figure 2.** Time evolution of: (a) the azimuthal mode number spectrum of the electrostatic potential in a logarithmic scale; (b) the empirical instantaneous growth rate of the kinetic energy of the background flow,  $\gamma_{m=0,n=0} = \partial_t \ln E_{0,0}$ .

is its singular value, that is, the weight that defines the relative importance [10] of the mode's contribution to  $\phi(r, \theta, t)$ .




**Figure 3.** (Left panel) The spatial character of the nine SVD modes that have the largest singular values. (Middle panel) The time evolution of the SVD modes, characterised by  $h_j(t)$  in equation (1). (Right panel) Singular values of the first twenty SVD modes, plotted on a logarithmic scale.

We emphasise that the functional form of each  $\Psi_j$  is entirely dictated *a posteriori* by the simulation outputs, without any prior assumptions. In appendix A, we outline how to perform the SVD method to the dataset observed in two-dimensional system. Figure 3 shows the dominant structure functions  $\Psi_j$ , and their time evolution  $h_j$ . These are the top nine modes, in descending order of their singular values  $s_j$ . We note that the two-dimensional pattern of  $\phi(r, \theta, t)$  dynamically changes in time, as shown in figure 2 for Fourier space, or figure 1 of [17] for real space. The nine spatial structure functions shown in the left panel of figure 3 capture background profile deformation (modes 1 and 4) and turbulent structures and fluctuations (the remaining modes). Here the mode ID numbers 1, 2,  $\dots$ , 9 are in descending order of the singular values  $s_j$ . These are the dominant modes in this SVD, as can be seen from the right panel of figure 3 where values of  $s_j$  are plotted on a logarithmic scale for modes up to the hundredth. In the present study, we truncate the SVD modes at the fortieth, for which the energy transfer can be calculated within acceptable accuracy. Convergence checks in relation to mode truncation are discussed in appendix B.

One can see from figure 3 that there are several paired modes whose spatial structures are similar and whose singular

values are nearly the same, which differ only in their phase. Examples of paired modes are: ID = 2, 3, ID = 5, 6, and ID = 7, 8, 9. These are complex conjugate pairs, which in combination represent the azimuthal propagation, akin to the combination of sine and cosine components in Fourier mode decomposition. Thus, the summation of the pair should be treated as a single structure. There are also unpaired modes, for example modes ID = 1 and ID = 4, which are almost azimuthally homogeneous and do not propagate in the azimuthal direction. The pairing of modes can be demonstrated from their envelopes calculated from  $h_j$  as shown in the middle panel of figure 3. The envelopes, delineated in black, change slowly in time; here the envelopes represent  $\sqrt{h_2^2 + h_3^2}$  (for ID = 2, 3),  $\sqrt{h_5^2 + h_6^2}$  (for ID = 5, 6) and  $\sqrt{h_7^2 + h_8^2 + h_9^2}$  (ID = 7, 8, 9). One can see that the temporal behaviours of modes with ID = 5–9 are intermittent (time scales of their growth and decay are comparable to their frequency), while the modes with ID = 2–3 are coherent. We wish to capture low-dimensional phenomenology wherever possible, and therefore choose to reduce the number of degrees of freedom by taking a summation from mode ID = 5 to ID = 40; we then treat the resulting summed entity as a single structure.





**Figure 4.** Dominant turbulent structures extracted by summation and truncation of the modes in figure 3. Mode A is invariant against azimuthal rotation, and the dominant component of Mode B is an azimuthally symmetric zonal pattern. Modes C and D are the turbulent structures.

The spatiotemporal behaviour of the electrostatic potential is thus decomposed into four structures,  $\phi_A, \phi_B, \phi_C, \phi_D$ , which are defined as follows:

$$\phi(r, \theta, t) \approx \sum_{\zeta=A \sim D} \phi_{\zeta}(r, \theta, t), \quad (2a)$$

$$\phi_A(r, \theta, t) = \varphi_1(r, \theta, t), \quad (2b)$$

$$\phi_B(r, \theta, t) = \varphi_4(r, \theta, t) \quad (2c)$$

$$\phi_C(r, \theta, t) = \varphi_2(r, \theta, t) + \varphi_3(r, \theta, t), \quad (2d)$$

$$\phi_D(r, \theta, t) = \sum_{j=5}^{40} \varphi_j(r, \theta, t), \quad (2e)$$

where  $\varphi_j(r, \theta, t) = s_j \Psi_j(r, \theta) h_j(t)$ . Snapshots of each of these four structures at  $t = 3200$  are shown in figure 4. Mode A corresponds to the background potential deformation; mode B to the zonal flow; mode C to the coherent KH mode; and mode D to the intermittent structure. Here, we note mode A and mode B are distinguished from their radial wavenumber;  $k_r \leq \pi/a$  (mode A) and  $k_r > \pi/a$  (mode B), where  $k_r$  is the

radial wavenumber of the electrostatic potential with  $m = n = 0$ . We refer to these as extracted structures of the electrostatic potential. The kinetic energy of each structure,  $E_{\zeta}$  can be evaluated from the definition


$$E_{\zeta} = \frac{1}{2} \int |\nabla_{\perp} \phi_{\zeta}|^2 d^3x, (\zeta = A \sim D). \quad (3)$$

The time evolution of the kinetic energy of each structure is illustrated in figure 5(a). Information about the relative phase of the time-evolving kinetic energy in pairs of modes can also be inferred from the phase plots in the right panel of figure 5. The process for constructing these plots is identical to that for Lissajous figures. While, strictly, the latter term applies to superimposed simple harmonic motions, we use it more colloquially here and in figures 8 and 9. Lissajous figures (in this sense) are widely used to characterise strongly non-linear structures in plasmas, see for example the treatment of heat pulse propagation in the LHD heliotron-stellarator in figures 1, 3, 5, 6 and 7 of [33] and figure 7 of [34].

The behaviour of the energy that is evident in figure 5 is similar to that extracted by using the DMD method, see in particular figures 4 of [17]. The background potential (mode A) and the coherent KH mode C compete with each other. When the magnitude of mode C increases to a certain threshold, the zonal flow mode B is excited together with the intermittent structure mode D. After the appearance of modes B and D, mode C decreases and mode A increases. This cycle then repeats—it is an LCO. The period of the LCO is synchronised with the zonal flow oscillation. The time evolution of the local value of the zonal potential (mode B) at different radial locations is plotted in figure 5(b). The phase of mode B at  $r/a = 0.23$  is the reverse of the phase at  $r/a = 0.6$ . The plot of the time-evolution of the potential at these two radial locations exhibits standing-wave-like structure. Successive zero-crossings—that is, phase reversal events—are synchronous with successive cycles of the LCO, where the frequencies of the LCO and the zonal flow are similar and the phase relation between the zonal flow and the LCO does not vary over time. The zonal flow mode B always peaks sharply just before each abrupt increase of the background flow mode A, and the phase of mode B reverses after the burst of mode A.

#### 4. Energy transfer dynamics among SVD modes

The energy transfer among the structures extracted in the preceding section can be characterised in terms of an energy transfer function that is based on the vorticity equation. The kinetic energy evolution equation can be derived by multiplying both sides of the vorticity equation by  $\phi$  and performing a spatial integration. The details of the derivation is described in appendix C. Here,  $\phi$  is decomposed into the SVD structures from mode A to mode D as in equation (2). The kinetic energy of Mode  $\zeta$  can be written as ( $\zeta = A \sim D$ )



**Figure 5.** Characterising limit cycle oscillations. Left panel. Time evolution of: (a) the kinetic energy of each of the four structures A to D identified in figure 4; (b) the local values of the zonal flow potential (Mode (B) at widely separated radial locations  $r/a = 0.23$  and  $0.6$ , at angular position  $\theta = 0$ . Right panel. Three Lissajous figures plotting the time evolution of the kinetic energy of the three Modes A, B and D against that of Mode C, during the time interval  $3080 < t < 3250$ . The arrows show the direction of time evolution.

$$\partial_t E_\zeta = \sum_{\alpha, \beta=A \sim D} J(\zeta|\alpha, \beta) - 2\nu E_\zeta - \int \phi_\zeta S d^3x + F_\zeta, \quad (4a)$$

$$J(\zeta|\alpha, \beta) = \int \phi_\zeta [\phi_\alpha, \nabla_\perp^2 \phi_\beta] d^3x, \quad (4b)$$


where the square brackets in equation (4b) denote the convective derivative defined as  $[\phi, \nabla_\perp^2 \phi] = r^{-1} (\partial_r \phi \partial_\theta \nabla_\perp^2 \phi - \partial_r \nabla_\perp^2 \phi \partial_\theta \phi)$ , and  $\nu$  and  $S$  are the ion-neutral collision frequency and the vorticity source, respectively. The term  $F_\zeta$  is the other nonlinearities introduced in appendix C. In this study, we focus on the term related to the convective non-linearity. The energy transfer function,  $J(\zeta|\alpha, \beta)$ , quantifies the energy flow to the mode labelled  $\zeta$  that arises from non-linear coupling between the modes labelled  $\alpha$  and  $\beta$ . Positive (or negative)  $J(\zeta|\alpha, \beta)$  contributes to driving (or damping). The terms in the RHS of equation (4a) are linear, except for the energy transfer term. In the present SVD context, equation (4a) can be viewed as a natural extension of the energy transfer equation in Fourier space [35, 36]. It is well adapted to understanding the energy interaction among the turbulent structures identified using SVD, which contain multiple Fourier components. In the absence of such structures, a Fourier-based approach to energy transfer would suffice for zonal flows, which can be described by a single Fourier component. It follows that the evidently non-linear features of the system dynamics, for example those shown in figure 5 must stem from the energy transfer among the turbulent structures. In order to understand the underlying physics of the system, it is therefore necessary to examine the behaviour of the energy transfer function in greater detail. As a preliminary, we note that  $J(\zeta|\alpha, \beta)$  is not symmetric with respect to  $\alpha$  and  $\beta$ . Given the four structures A to D, there are thus sixteen relevant mode pairings. Table 1 specifies how we label (‘Coupling ID’) these pairings using the first sixteen integers. We note that the SVD

**Table 1.** List of coupling IDs. Each number denotes the combination of structures which contribute to the coupling.

Coupling ID	Combination of structures
1	A + A
2	A + B
3	A + C
4	A + D
5	B + A
6	B + B
7	B + C
8	B + D
9	C + A
10	C + B
11	C + C
12	C + D
13	D + A
14	D + B
15	D + C
16	D + D

method does not involve well-defined matching criteria with respect to frequency or wavenumber because, in general, SVD modes contain multiple Fourier modes. In the present paper, mode D is a good example of this.

Figure 6 seeks to capture the temporal variation of  $J(\zeta|\alpha, \beta)$  for each structure A to D, resulting from each of the sixteen coupling combinations, from two different perspectives. The four left panels, figure 6(a)–(d), plot the instantaneous values that are taken by each  $J(\zeta|\alpha, \beta)$  at each unit time-step during the interval  $3000 < t < 4000$  which was also considered in figures 2 and 5. For example, figure 6(c) shows that for mode C, the four sources/sinks of non-linear drive that contribute most strongly to energy transfer (both positive and negative) are the mode pairings with coupling ID = 4, 8, 13 and 14. From table 1 in  $(\alpha, \beta)$  terms, these pairings are (A,D),



**Figure 6.** (a)–(d): Time variation of the energy transfer functions for each structure. (e)–(h): Temporal evolution of the energy transfer functions for each structure, where (e)  $J(A|\alpha, \beta)$ , (f)  $J(B|\alpha, \beta)$ , (g)  $J(C|\alpha, \beta)$ , and (h)  $J(D|\alpha, \beta)$ , where  $\alpha + \beta$  denotes the coupling between mode  $\alpha$  and mode  $\beta$ , in the order corresponding to equation (4b).


(B,D), (D,A) and (D,B). Conversely, from figure 6(b) it is evident that energy transfer to and from mode B is dominated by the mode pairing with coupling ID = 11, 12, 15, 16 that are  $(\alpha, \beta) = (C,C), (C,D), (D,C), (D,D)$ . From figure 6(a) one can see that the background flow, mode A, is affected primarily by the self-couplings (C,C) and (D,D) and by the pair couplings (C,D) and (D,C). There is no coupling between the zonal flow and the background flow, which is because the background flow does not generate the doppler shift for the zonal flow. The background and the zonal flows couple indirectly through the deformation of the fluctuations. We note from figure 4 that the azimuthally symmetric background flow is contained in mode A and partly in mode B. This flow is driven by the vorticity source. It follows that net negative non-linear energy transfer to mode A acts to relax the background flow. In contrast, for mode C and mode D, non-linear damping due to self-coupling is found to be weak, in that the contributions from the coupling ID = 11 for mode C and 16 for mode D are small. The four right panels, figures 6(e)–(h) plot the explicit time evolution of  $J(\zeta|\alpha, \beta)$  on a colour scale, for the four modes (one panel per mode) in relation to the 16 mode pairings. The energy transfers among the structures are strongly non-time-stationary: the discrete bursts of activity correlate necessarily with those in figure 5 which align with the LCO period, as discussed previously. In the following, in order to reduce the number of coupling combinations, we treat the couplings of mode- $\alpha$  with mode- $\beta$  and of mode- $\beta$  with mode- $\alpha$  together as  $J(\zeta|\alpha, \beta) + J(\zeta|\beta, \alpha) \rightarrow J(\zeta|\alpha, \beta)$ .

The energy transfer functions for each structure during a single LCO period,  $3120 < t < 3220$ , are illustrated in figure 7 where only the most important couplings are shown. There are


three phases, because the energy flow pattern changes significantly between the first half ( $3120 < t < 3180$ ), the middle regime ( $t \sim 3190$ ), and the second half ( $3200 < t < 3220$ ). This pattern corresponds to the growth, saturation and damping of the zonal flow, mode B. The energy transfer to the zonal flow changes its sign at  $t \sim 3190$ , at which time the zonal flow peaks in energy. This saturation is caused by the reversal in energy transfer  $J(B|C, D)$ . This implies the coupling  $C + D$  is important for the non-linear saturation of the zonal flow. Simultaneously, there is an abrupt onset to the non-linear drive of the background flow (mode A), due to the non-linearity of mode D, as shown in figure 7(b), see figure 7(a); the energy of mode A, previously decaying, starts to increase.

The  $x$ -axis of all three panels of figure 8 is  $J(C|A, C)$ . This is chosen because, in general,  $J(\zeta|A, \zeta)$  represents the generalised linear drive of the mode labelled  $\zeta$  by the KH instability, which arises from deformation of the background potential mode A. This drive would be linear, notwithstanding its apparently non-linear mathematical character, in an ordering scheme where mode A is zeroth order while the other modes are order-epsilon. In figure 8 the focus is thus on the different non-linear drives of the coherent KH mode C, in relation to the linear drive. The linear drive for the coherent KH mode C, which corresponds to  $J(C|A, C)$ , changes its role in accordance with the phase of the LCO. It is apparent from figure 7(a) that, during the first half of the LCO, mode C obtains energy from its coupling with the background potential deformation mode A; that is, mode C is linearly destabilized by the flow inhomogeneity. This phase of the LCO ends at  $t \sim 3185$  in figure 7 and this determines the parameter  $t_2$  which is used throughout figures 8 and 9 to denote the end of the first half (blue trace)





**Figure 7.** Time evolution of (a) kinetic energy and (b)–(e) energy transfer function for each structure. Only the energy transfer functions for the most important non-linear couplings are shown.




**Figure 8.** Lissajous figures for the three dominant energy transfer functions for the KH mode C,  $J(C|\alpha, \beta)$ , where the arrows show the direction of the time-evolution. Here,  $t_1 = 3080$ ,  $t_2 = 3185$ ,  $t_3 = 3195$  and  $t_4 = 3250$ .

of the LCO cycle. Thereafter, figure 8(a) shows that coupling  $J(C|B, C)$  of the KH mode C with the zonal flow mode B acts to suppress mode C; this is a back reaction which drives the zonal

flow. In the Lissajous figures 8 and 9 the system trajectory during the three time intervals  $3080 < t < 3185$ ,  $3185 < t < 3195$  and  $3195 < t < 3250$  is plotted using blue, red and black lines, respectively. The non-linear effect of the intermittent structure mode D, characterised by  $J(C|B, D)$  in figure 8(b) has a negligible role during the first half of the LCO. It begins to acquire a small negative value when the linear drive of mode C,  $J(C|A, C)$ , approaches its maximum value of 0.015, and then rises abruptly after the onset of the zonal flow (red trace). The plot of  $J(C|D, D)$  in figure 8(c) shows that the non-linear self-interaction of mode D has a negative effect on mode C in the first and middle phase of the LCO. At the conclusion of the zonal flow drive, as  $J(C|A, C)$  declines towards zero,  $J(C|D, D)$  starts (black trace) to contribute to the drive of mode C. In this way, the non-linear modifications to mode C,  $J(C|B, D)$  and  $J(C|D, D)$ , change their character after the onset of the zonal flow.

The linear drive for the intermittent structure mode D corresponds to  $J(D|A, D)$ . The Lissajous figures for  $J(D|\alpha, \beta)$  during the LCO are shown in figure 9 and demonstrate that  $J(D|A, D)$  changes its role in accordance with the phase of the LCO, like  $J(C|A, C)$  in figure 8 [17]. Figure 9(a) shows that for mode D, unlike for mode C, the deformation of the KH mode C due to its non-linear coupling with the background flow mode A contributes to the drive. Deformation of mode C by the zonal flow B also affects the energy transfer to Mode D,



**Figure 9.** Lissajous figures for the four dominant energy transfer function for the intermittent mode D,  $J(D|\alpha, \beta)$ , where the arrows show the direction of the evolution. Here,  $t_1 = 3080$ ,  $t_2 = 3185$ ,  $t_3 = 3195$  and  $t_4 = 3250$ .


as can be inferred from figure 9(b). This shows a rather abrupt reversal between the first half (blue trace) and second half (red, then black, traces) of the LCO, giving rise to a well-defined figure 8. The non-linear coupling of Mode D with the zonal flow mode B is shown in figure 9(c). The sign of  $J(D|B, D)$  is always negative, implying that it contributes to the suppression of mode D throughout the LCO. As seen in the Lissajous figure for  $J(D|C, D)$  in figure 9(d) non-linear coupling with mode C drives mode D during the first half (blue trace) of the LCO; damps it during the middle phase (red trace) and much of the final phase (black trace).

The energy flow patterns inferred from figures 8 and 9 during a single LCO period are summarized in figure 10. We have found that important roles in the sustainment of the LCO are played by the secondary structures, namely the zonal flow mode B and the intermittent structure mode D, notably through their non-linear coupling with the KH mode C.

We now perform a quantitative check on our inference that the abrupt energy input to the background flow mode A at  $3185 < t < 3195$  causes the abrupt increase in the background flow that is shown in figure 2. Dividing both sides of equation (4a) by  $E_C$ , an equation for the time constant of mode A,  $\gamma_A$ , can be formally derived as

$$\gamma_A = E_A^{-1} \sum_{\alpha, \beta} J(A|\alpha, \beta) - \nu. \quad (5)$$

From figure 2 the time constant at the abrupt increase can be estimated as  $\gamma_A \sim 2 \times 10^{-2} \gg \nu = 5 \times 10^{-3}$ . This value can be explained as  $E_A^{-1} \sum_{\alpha, \beta} J(A|\alpha, \beta) \sim 2 \times 10^{-2}$ , where we use  $E_A = 0.5$  and  $\sum_{\alpha, \beta} J(A|\alpha, \beta) \approx J(A|D, D) \sim 0.01$ . This tends to confirm that the abrupt increase of the background flow mode A is realised by the non-linear self-coupling of the nonlinearly driven intermittent structure D. We note that a strong role for the zonal flow mode B in the abrupt change of the background flow is also implicit. This is because, as we



**Figure 10.** Schematic view of the energy flow among turbulent structures during the limit cycle oscillation, where A–D in the circles denote the Mode A–D, respectively, and the arrows show the directions of the energy flow. The pattern of the energy flow changes during one period of the LCO.

have already shown, deformation of modes C and D by mode B abruptly alters the characteristics of non-linear energy transfer involving modes C and D.

## 5. Discussion and Conclusions

Using the SVD method, the global phenomenology of the simulated turbulent plasma has been converted into a low-dimensional dynamical system. The electrostatic potential is decomposed into four elements, which are apparently the minimal set: the deformation of the background; the zonal flow; the coherent KH mode; and the intermittent structure. The kinetic energy content of these four modes is found to exhibit an LCO, in which an abrupt change of the background potential is synchronized with the zonal flow period.

Wider application of the SVD method, as presented here, to experimental data from plasmas should be relatively straightforward. While there is no constraint on the number of measurement points in space and time, we note that simultaneous measurements are necessary to construct the spatiotemporal information matrix,  $\Phi(x_i, t_j)$ , which is introduced in appendix A. Many typical plasma diagnostics, such as fast camera imaging, Langmuir probes, reflectometers, heavy ion beam probes, beam emission spectroscopy, and tomography, yield datasets which appear suitable for the SVD method. Turning specifically to cylindrical plasmas of the kind considered here, we anticipate that the four key structures identified in the present paper will again be found in future experiments.

The SVD approach has enabled us to quantify the multiple non-linear energy flows that underly this LCO. We first construct and evaluate the energy transfer function for each

turbulence structure, based on a vorticity equation. This then provides physical understanding of how the LCO is sustained by dynamical changes in the energy transfer among structures over the LCO period. In summary, coupling between the coherent mode and the intermittent structure drives the zonal flow, which provides resilience for the background potential, which causes the abrupt deformation of the background.

In some previous theoretical approaches, the LCO considered here is often described in terms of a predator–prey model [37] that involves only three elements, namely the background, the zonal flow and microturbulence. Thus, the results in the present paper imply that this conventional model is incomplete. Furthermore, we have found that the hitherto neglected intermittent structure plays crucial roles in the dynamics of the system. The important role of the intermittent structure in the abrupt change in transport has also been identified experimentally [38].

In addition, we have found that the zonal flow oscillation is synchronised with the LCO. The physical mechanism that determines the frequency of the zonal flow remains an unresolved theoretical issue. In the context of the present work, we note that in basic experiments, the excitation of an intermittent structure, referred to as ‘splash’, has been observed to be synchronised with the zonal flow oscillation [39, 40]. This tendency is similar to the behaviour of the intermittent mode D in the present study. This motivates our conjecture that the generation and annihilation of the intermittent structure, together with the deformation of the background flow, may determine the frequency of the zonal flow.

In conclusion, the SVD approach has provided significant new physical insights into the phenomenology of this classical non-linear KH plasma system.

## Acknowledgments

We acknowledge useful discussions with S Maeyama of Nagoya University and with O Samant of CFSA, Warwick University. This work was partly supported by a grant-in-aid for scientific research of JSPS, Japan (16K18335, 17H06089, 16H02442, 15H02155, 15H02335, 18K03578), the collaboration programs of NIFS (NIFS18KNST137, NIFS19KNST151), Asada Science Foundation, Progress 100 of Kyushu University (NB80645028), and the RIAM of Kyushu University. This work was carried out partly within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014–2018 and 2019–2020 under Grant Agreement No 633053. The work received support from the RCUK Energy Programme grant no. EP/T012250/1. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

## Appendix A. SVD analysis in two-dimensional system

Here we briefly explain how to perform SVD analysis to a set of spatio-temporal data in two-dimensional system.

The spatio-temporal data in 2D system can be obtained in a three-dimensional array,  $\phi(x_1, x_2, t)$ , where  $x_j$  is the spatial coordinate. The size of this array is  $M_1 \times M_2 \times N$ , where  $M_j$  and  $N$  are the number of measurement points regarding  $x_j$  and time, respectively. In order to perform the SVD, the three-dimensional array is converted to two-dimensional array,  $\phi(x_1, x_2, t) \rightarrow \Phi(X, t)$ , where the size of the matrix is  $M \times N$ , where  $M = M_1 M_2$ . Then, the spatio-temporal information matrix, whose component is  $\Phi(X_i, t_j) = \Phi_{ij}$ , can be generally decomposed into  $\Phi_{ij} = U \Sigma V^T$ , where  $U$  and  $V$  are  $M$ th and  $N$ th order orthogonal matrix, and  $\Sigma$  is  $M \times N$  diagonal matrix. This expression can be rewritten by

$$\Phi = \sum_{i=1}^p s_i \mathbf{u}_i \mathbf{v}_i^T, \quad (\text{A1})$$

where  $\mathbf{u}_i$  and  $\mathbf{v}_i$  are the column vector of  $U$  and  $V$ , respectively, and  $s_i$  is the component of  $\Sigma$ . The vectors  $\mathbf{u}_i$  and  $\mathbf{v}_i$  correspond to spatial and temporal structures, respectively. Then, one can obtain the 2D pattern by re-converting  $X \rightarrow (x_1, x_2)$ .

Care is needed when spatiotemporal data, expressed with respect to cylindrical coordinates, is decomposed into SVD modes. At first sight, it appears that the orthogonality condition for SVD modes should be given by

$$\int \phi_\zeta(r, \theta) \phi_{\zeta'}(r, \theta) dr d\theta = \delta_{\zeta, \zeta'}, \quad (\text{A2})$$

where  $\phi_\zeta$  denotes the SVD mode that has mode ID  $\zeta$ . We note that there is no apparent requirement for a Jacobian  $r$  in the integration with respect to the independent variables  $r$  and  $\theta$  in equation (A2). However, when analysing the energy transfer, the spatial integral  $\int \dots r dr d\theta$  is performed. This is necessary for computational accuracy in the context of an intrinsically cylindrical plasma system. It then appears that, for consistency, the SVD orthogonality condition also needs to incorporate a Jacobian  $r$ , whereas equation (A2) does not.


This conundrum is resolved by defining SVD orthogonality with respect to a Cartesian coordinate system  $(x, y)$ . On transforming to cylindrical coordinates, this SVD orthogonality condition becomes

$$\int \phi_\zeta(r, \theta) \phi_{\zeta'}(r, \theta) r dr d\theta = \delta_{\zeta, \zeta'}. \quad (\text{A3})$$

Instead of equation (A2). In the present study, we compute the SVD in Cartesian coordinates, after transforming  $\phi(r, \theta, t)$  to  $\phi(x, y, t)$ . This implicitly guarantees the orthogonality condition equation (A3) and is consistent with energy transfer calculations that are carried out exclusively in the more natural  $(r, \theta, t)$  coordinates.

## Appendix B. Convergence check for mode truncation

Here, we check the convergence for the truncation of the SVD modes. The number of SVD modes generated by the SVD process is equal to the number of time steps. In the present study,



**Figure B1.** Difference between the input data and the reconstructed data, plotted against the SVD mode number  $M$  at which the SVD decomposition is truncated: (a) for spatial structures; (b) for spatially integrated variables.

we address the data from  $t = 3000$  to  $t = 4000$  with  $\Delta t = 1$ , where  $\Delta t$  is the time step, so that one thousand SVD modes are produced. This is clearly excessive in relation to the objective of capturing low-dimensional dynamics, which provides the rationale for our use of SVD: truncation is therefore necessary. Importantly, the calculation of spatial derivatives of the SVD-decomposed data is central to our analysis of the non-linear physics. It follows that mode truncation must be robustly convergent in two senses: in reconstructing the original data; and in relation to spatial derivative variables. We note that in order to capture physical effects that involve higher order spatial derivatives, the retention of finer structure is necessary. Conversely, for spatially integrated quantities, convergence under truncation is faster than for spatial structures and their derivatives.

Let us denote the reconstructed data under truncation at the  $M$ th SVD mode as

$$\phi^{(M)} = \sum_{j=1}^M s_j \Psi_j h_j. \quad (\text{B1})$$

We calculate the turbulence stress  $\Pi_{r,\theta}^{(M)}$  (which requires a first order spatial derivative) and the vorticity  $\Omega^{(M)}$  (involving the second derivative), using the reconstructed data  $\phi^{(M)}$ ,

$$\Pi_{r,\theta}^{(M)} = -\partial_r \phi^{(M)} \frac{1}{r} \partial_\theta \phi^{(M)}, \quad (\text{B2})$$

$$\Omega^{(M)} = \nabla_\perp^2 \phi^{(M)}. \quad (\text{B3})$$

We then compare these with the corresponding quantities calculated from the original data  $\phi$ . Their deviations are evaluated from

$$\epsilon_X^{(M)} = \sqrt{\frac{\langle \int |X^{(M)} - X|^2 dx^3 \rangle_t}{\langle \int |X|^2 dx^3 \rangle_t}}. \quad (\text{B4})$$

Here  $X$  corresponds to  $\phi, \Pi_{r,\theta}, \Omega$  calculated from the original data, whereas  $X^{(M)}$  is obtained under SVD mode truncation at

mode  $M$ , as in equations (B1) to (B3) and  $\langle \dots \rangle_t$  denotes the time average. As shown in figure B1(a) the variables with the higher order derivatives converge more slowly.

Next, we check the convergence of the variables which involve spatial integration: the kinetic energy of the intermittent structure (mode D) and the energy transfer function between the background flow (mode A) and mode D. The spatio-temporal data of mode D under the mode truncation is introduced as

$$\phi_D^{(M)} = \sum_{j=5}^M s_j \Psi_j h_j. \quad (\text{B5})$$

The kinetic energy of mode D under truncation is denoted by  $K_D^{(M)}$ , and we denote the energy transfer to the background flow due to the self-non-linearity of mode D, under truncation, by  $J^{(M)}(A|D, D)$ . These are given by

$$K_D^{(M)} = \frac{1}{2} \int |\nabla_\perp \phi_D^{(M)}|^2 d^3x, \quad (\text{B6})$$

$$J^{(M)}(A|D, D) = \int \phi_A [\phi_D^{(M)}, \nabla_\perp^2 \phi_D^{(M)}] d^3x. \quad (\text{B7})$$

The deviation of these quantities from those calculated from the original data is defined as

$$\delta_Y^{(D)}(M) = \sqrt{\frac{\langle |Y^{(M)} - Y|^2 \rangle_t}{\langle |Y|^2 \rangle_t}}, \quad (\text{B8})$$

where  $Y$  corresponds to  $K_D$  and  $J(A|D, D)$ . The deviation  $\delta_Y^{(D)}$  is plotted in figure B1(b). This shows that, empirically, their convergence is faster than for the spatial variables with similar trends, even though the energy transfer function has a higher order derivative (third order). In the present study, we therefore truncate the SVD mode decomposition at  $M = 40$  in order to retain 1% accuracy for the spatially integrated variables.

### Appendix C. Derivation of full set of energy equations

The energy equations for the SVD modes are systematically derived in this appendix. The basic equations used in this simulation are given as [28]

$$\partial_t N = -[\phi, N] - D \nabla_{\parallel}^2 (\phi - N) + \mu_N \nabla_{\perp}^2 N + S_N, \quad (C1)$$

$$\begin{aligned} \partial_t \nabla_{\perp}^2 \phi &= -[\phi, \nabla_{\perp}^2 \phi] - D \nabla_{\parallel}^2 (\phi - N) - \nu_{in} \nabla_{\perp}^2 \phi \\ &\quad - \nu_{in} \nabla N \cdot \nabla_{\perp} \phi \\ &\quad - \nabla N \cdot d_t \nabla_{\perp} \phi + \mu_U \nabla_{\perp}^4 \phi + S_U, \end{aligned} \quad (C2)$$

where  $D$  is the electron parallel diffusion,  $\mu_N$  and  $\mu_U$  are the diffusion coefficients for particle and vorticity, and  $\nu_{in}$  is the ion-neutral collision frequency. The particle and vorticity sources are given by  $S_N$  and  $S_U$ . Here, we decompose the fields,  $N$  and  $\phi$ , with the SVD modes as

$$N = \sum_{\zeta} N_{\zeta}, \quad (C3)$$

$$\phi = \sum_{\zeta} \phi_{\zeta}, \quad (C4)$$

where  $N_{\zeta}$  and  $\phi_{\zeta}$  are the SVD modes for density and potential. It is noted that each SVD mode satisfies the orthogonality condition as

$$\int \phi_{\zeta} \phi_{\zeta'} dx^3 = \int N_{\zeta} N_{\zeta'} dx^3 = \delta_{\zeta, \zeta'}, \quad (C5)$$

where  $\zeta'$  is the Kronecker delta. Multiplying  $N_{\zeta}$  and  $\phi_{\zeta}$  to density and vorticity equation, respectively, and performing the spatial integrals, the following energy equations are obtained. In the derivation, the orthogonality condition is used.

$$\begin{aligned} \partial_t \int \frac{1}{2} N_{\zeta}^2 dx^3 &= - \sum_{\alpha, \beta} \int N_{\zeta} [\phi_{\alpha}, N_{\beta}] dx^3 + \int N_{\zeta} S_N dx^3 \\ &\quad - D \sum_{\zeta'} \int N_{\zeta} \nabla_{\parallel}^2 (\phi_{\zeta'} - N_{\zeta'}) \\ &\quad - \mu_N \int |\nabla_{\perp} N_{\zeta}|^2 dx^3 - \sum_{\zeta \neq \zeta'} \int N_{\zeta} \partial_t N_{\zeta'}, \end{aligned} \quad (C6)$$

$$\begin{aligned} \partial_t \int \frac{1}{2} |\nabla_{\perp} \phi_{\zeta}|^2 dx^3 &= \sum_{\alpha, \beta} \int \phi_{\zeta} [\phi_{\alpha}, \nabla_{\perp} \phi_{\beta}] \\ &\quad - \nu_{in} \int |\nabla_{\perp} \phi_{\zeta}|^2 dx^3 \\ &\quad - \int \phi_{\zeta} S_U dx^3 + F_{\zeta}, \end{aligned} \quad (C7)$$

where  $F_{\zeta}$  is given as

$$\begin{aligned} F_{\zeta} &= \sum_{\alpha, \beta} \int \phi_{\zeta} [\nu_{in} \nabla N_{\alpha} \cdot \nabla_{\perp} \phi_{\beta} + \nabla N_{\alpha} \cdot d_t \nabla_{\perp} \phi_{\beta}] dx^3 \\ &\quad + \sum_{\zeta'} \int \phi_{\zeta} [D \nabla_{\parallel}^2 (\phi_{\zeta'} - N_{\zeta'}) - \mu_U \nabla_{\perp}^4 \phi_{\zeta'}] dx^3 \\ &\quad + \sum_{\zeta \neq \zeta'} \int \nabla_{\perp} \phi_{\zeta} \cdot \partial_t \nabla_{\perp} \phi_{\zeta'} dx^3. \end{aligned} \quad (C8)$$

These are the full set of energy equation. In this study, only the potential is decomposed with the SVD, and we focus on the convective non-linearity. The full analysis using the multi-field SVD analysis for the density and potential could be future work.

### ORCID iDs

M Sasaki  <https://orcid.org/0000-0001-6835-1569>  
 T Kobayashi  <https://orcid.org/0000-0001-5669-1937>  
 Y Kawachi  <https://orcid.org/0000-0002-5222-6082>  
 H Arakawa  <https://orcid.org/0000-0001-9793-099X>

### References

- [1] Itoh S-I 2009 Structure formation in turbulent plasmas *Plasma Fusion Res.* **4** 038–038
- [2] Diamond P H, Itoh S I, Itoh K and Hahn T S 2005 Zonal flows in plasma—a review *Plasma Phys. Control. Fusion* **47** R35
- [3] Gürçan OD, Diamond P H, Hahn T S and Singh R 2007 Intrinsic rotation and electric field shear *Phys. Plasmas* **14** 042306
- [4] Litaudon X 2006 Internal transport barriers: critical physics issues? *Plasma Phys. Control. Fusion* **48** A1
- [5] Wagner F 2007 A quarter-century of h-mode studies *Plasma Phys. Control. Fusion* **49** B1
- [6] Biglari H, Diamond P H and Terry P W 1990 Influence of sheared poloidal rotation on edge turbulence *Phys. Fluids B: Plasma Phys.* **2** 1–4
- [7] Imamura H 2006 Introduction to nonlinear analysis for engineerings, suurikougakusha-sha co.,ltd *Appl. Math.* **16** 185–6
- [8] Miki K, Diamond P H, Hahn S-H, Xiao W W, Gürçan Ozgür D and Tynan G R 2013 Physics of stimulated l→h transitions *Phys. Rev. Lett.* **110** 195002
- [9] Sasaki M, Itoh K, Itoh S-I and Kasuya N 2012 Zonal flows induced by symmetry breaking with existence of geodesic acoustic modes *Nucl. Fusion* **52** 023009
- [10] Dudok de Wit T, Pecquet A-L, Vallet J-C and Lima R 1994 The biorthogonal decomposition as a tool for investigating fluctuations in plasmas *Phys. Plasmas* **1** 3288–300
- [11] Hatch D R, Poeschel M J, Jenko F, Nevins W M, Terry P W and Doerk H 2013 Magnetic stochasticity and transport due to nonlinearly excited subdominant microtearing modes *Phys. Plasmas* **20** 012307
- [12] Ph Van Milligen B, Sánchez E, Alonso A, Pedrosa M A, Hidalgo C, Martín De Aguilera A and López Fraguas A 2014 The use of the biorthogonal decomposition for the identification of zonal flows at tj-ii *Plasma Phys. Control. Fusion* **57** 025005
- [13] Ohdachi S, Toi K and Fuchs G TEXTOR team, LHD Experimental Group et al 2006 Magnetic islands observed by a fast-framing tangentially viewing soft x-ray camera on lhd and textor *Plasma Sci. Technol.* **8** 45



- [14] Marchetto C, Alessi E, Boncagni L, Galperti C, Nowak S and Sozzi C 2012 Application of SVD algorithm to a set of real time Mirnov coil signals in FTU tokamak *Conf. on Plasma Phys.* pp 2–64
- [15] Gerksic S *et al* SVD-based fast online MPC for iter plasma current and shape control
- [16] Athanasiu M A and Pavlos G P 2001 SVD analysis of the magnetospheric AE index time series and comparison with low-dimensional chaotic dynamics *Nonlinear Process. Geophys.* **8** 95–125
- [17] Sasaki M, Kawachi Y, Dendy R O, Arakawa H, Kasuya N, Kin F, Yamasaki K and Inagaki S 2019 Using dynamical mode decomposition to extract the limit cycle dynamics of modulated turbulence in a plasma simulation *Plasma Phys. Control. Fusion* **61** 112001
- [18] Kusaba A, Kuboyama T and Inagaki S 2020 Sparsity-promoting dynamic mode decomposition of plasma turbulence *Plasma Fusion Res.* **15** 1301001–1301001
- [19] Kaptanoglu A A, Morgan K D, Hansen C J and Brunton S L 2020 Characterizing magnetized plasmas with dynamic mode decomposition *Phys. Plasmas* **27** 032108
- [20] del Castillo-Negrete D, Hirshman S P, Spong D A and D’Azevedo E F 2007 Compression of magnetohydrodynamic simulation data using singular value decomposition *J. Comput. Phys.* **222** 265–86
- [21] Hatch D R, Jenko F, Bañón Navarro A, Bratanov V, Terry P W and Pueschel M J 2016 Linear signatures in nonlinear gyrokinetics: interpreting turbulence with pseudospectra *New J. Phys.* **18** 075018
- [22] Kuramoto Y 2003 *Chemical Oscillations, Waves and Turbulence* (New York: Dover)
- [23] Fujisawa A *et al* 1998 Discovery of electric pulsation in a toroidal helical plasma *Phys. Rev. Lett.* **81** 2256
- [24] Kobayashi T *et al* 2013 Spatiotemporal structures of edge limit-cycle oscillation before l-to-h transition in the jft-2m tokamak *Phys. Rev. Lett.* **111** 035002
- [25] Schmitz L, Zeng L, Rhodes T L, Hillesheim J C, Doyle E J, Groebner R J, Peebles W A, Burrell K H and Wang G 2012 Role of zonal flow predator-prey oscillations in triggering the transition to h-mode confinement *Phys. Rev. Lett.* **108** 155002
- [26] Birkenmeier G *et al* 2016 Magnetic structure and frequency scaling of limit-cycle oscillations close to l-to h-mode transitions *Nucl. Fusion* **56** 086009
- [27] Itoh S-I, Itoh K, Fukuyama A and Miura Y 1991 Edge localized mode activity as a limit cycle in tokamak plasmas *Phys. Rev. Lett.* **67** 2485
- [28] Sasaki M, Camenen Y, Escarguel A, Inagaki S, Kasuya N, Itoh K and Kobayashi T 2019 Formation of spiral structures of turbulence driven by a strong rotation in magnetically cylindrical plasmas *Phys. Plasmas* **26** 042305
- [29] Hasegawa A and Wakatani M 1987 Self-organization of electrostatic turbulence in a cylindrical plasma *Phys. Rev. Lett.* **59** 1581
- [30] Horton W, Tajima T and Kamimura T 1987 Kelvin–Helmholtz instability and vortices in magnetized plasma *The Phys. Fluids* **30** 3485–95
- [31] Sasaki M *et al* 2017 Topological bifurcation of helical flows in magnetized plasmas with density gradient and parallel flow shear *Phys. Plasmas* **24** 112103
- [32] Itoh K, Itoh S-I, Kosuga Y, Lesur M and Ido T 2016 Onset condition of the subcritical geodesic acoustic mode instability in the presence of energetic-particle-driven geodesic acoustic mode *Plasma Phys. Rep.* **42** 418–23
- [33] Dendy R O, Chapman S C and Inagaki S 2013 Modelling the measured local time evolution of strongly nonlinear heat pulses in the large helical device *Plasma Phys. Control. Fusion* **55** 115009
- [34] Zhu H, Dendy R O, Chapman S C and Inagaki S 2015 A quantitative model for heat pulse propagation across large helical device plasmas *Phys. Plasmas* **22** 062308
- [35] Manz P, Ramisch M and Stroth U 2009 Physical mechanism behind zonal-flow generation in drift-wave turbulence *Phys. Rev. Lett.* **103** 165004
- [36] Nagashima Y *et al* 2009 Observation of the parametric-modulational instability between the drift-wave fluctuation and azimuthally symmetric sheared radial electric field oscillation in a cylindrical laboratory plasma *Phys. Plasmas* **16** 020706
- [37] Kim E-jin and Diamond P H 2003 Zonal flows and transient dynamics of the l- h transition *Phys. Rev. Lett.* **90** 185006
- [38] Kobayashi T *et al* 2015 Azimuthal inhomogeneity of turbulence structure and its impact on intermittent particle transport in linear magnetized plasmas *Phys. Plasmas* **22** 112301
- [39] Arakawa H *et al* 2016 Eddy, drift wave and zonal flow dynamics in a linear magnetized plasma *Sci. Rep.* **6** 33371
- [40] Arakawa H *et al* 2019 Roles of solitary eddy and splash in drift wave–zonal flow system in a linear magnetized plasma *Phys. Plasmas* **26** 052305
- [41] Kim J S, Edgell D H, Greene J M, Strait E J and Chance M S 1999 MHD mode identification of tokamak plasmas from Mirnov signals *Plasma Phys. Control. Fusion* **41** 1399
- [42] Hatch D R, Jenko F, Bañón Navarro A, Bratanov V, Terry P W and Pueschel M J 2016 Linear signatures in nonlinear gyrokinetics: interpreting turbulence with pseudospectra *New J. Phys.* **18** 075018