

UKAEA-CCFE-CP(23)27

O.P. Bardsley, T.C. Hender, B.C. Lyons, N.M.
Ferraro, S.N. Gerasimov, JET contributors

Extended-MHD simulations of shattered pellet injection into an ohmic JET plasma

This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the UKAEA Publications Officer, Culham Science Centre, Building K1/O/83, Abingdon, Oxfordshire, OX14 3DB, UK.

Enquiries about copyright and reproduction should in the first instance be addressed to the UKAEA Publications Officer, Culham Science Centre, Building K1/O/83 Abingdon, Oxfordshire, OX14 3DB, UK. The United Kingdom Atomic Energy Authority is the copyright holder.

The contents of this document and all other UKAEA Preprints, Reports and Conference Papers are available to view online free at scientific-publications.ukaea.uk/

Extended-MHD simulations of shattered pellet injection into an ohmic JET plasma

O.P. Bardsley, T.C. Hender, B.C. Lyons, N.M. Ferraro, S.N. Gerasimov, JET contributors

Extended-MHD simulations of shattered pellet injection into an Ohmic JET plasma

O. P. Bardsley¹, T. C. Hender¹, B. C. Lyons², N. M. Ferraro³, S. N. Gerasimov¹, JET contributors*

¹ *United Kingdom Atomic Energy Authority, Culham Centre for Fusion Energy, Culham Science Centre, Abingdon, Oxon OX14 3DB, UK*

² *General Atomics, San Diego, CA 92121, USA*

³ *Princeton Plasma Physics Laboratory, Princeton, NJ 08543-0451, USA*

* *See the author list of 'Overview of JET results for optimising ITER operation' by J. Mailloux et al. to be published in Nuclear Fusion Special issue: Overview and Summary Papers from the 28th Fusion Energy Conference (Nice, France, 10-15 May 2021)*

Abstract

Disruptions pose a serious challenge for future tokamak power plants; at the large stored thermal and magnetic energies of ITER and DEMO plasmas, the heat loads and electromagnetic forces during disruptions must be mitigated. With this objective, JET experiments employing impurity shattered pellet injection (SPI) have recently been undertaken.

We present numerical models of SPI disruptions in JET using M3D-C¹, an extended-MHD code developed for the study of non-linear transient events in tokamaks. Among its extensions are a neutral gas shielding model for the ablation of frozen pellets within the plasma, and the non-coronal equilibrium model KPRAD for tracking impurity charge states and radiation. With these, plasma cooling both before the thermal quench (when there is little MHD activity) and during it (when instability provokes rapid penetration of impurities and cooling of the plasma core) can be modelled.

Our results concern cooling of the Ohmic JET plasma #95149 by a small shattered neon-deuterium pellet, modelled in M3D-C¹ both with linearised stability analyses of axisymmetric simulations, and with a fully non-linear three-dimensional model. Comparisons are made with experimental measurements of temperature profiles and radiated power during the pre-thermal quench phase, and the timing and nature of thermal quench onset as the pellet impinges on rational surfaces and excites low- n MHD activity. This analysis qualifies the numerical approach, paving the way to simulations of higher-energy JET discharges and predictions for future devices.




Figure 1: Experiment summary. Images are from KL8-E8WA camera with a 527nm filter, radiation measurements from KB5V bolometer, and mid-plane electron temperature from KK3F ECE diagnostic.





Figure 2: Equilibrium flux surfaces (blue, separatrix in green) and pellet shard cloud used in M3D-C¹ simulations. Shards are red circles, with radius proportional to the physical size. Grey lines show pellet trajectories.

Experimental results


We consider JET shot 95149, a shattered pellet injection (SPI) into a healthy Ohmic L-mode plasma at 2MA/2T. The SPI is neon (0.11g) with a deuterium shell injected from above; the pellet diameter is 4mm, the smallest available, but this is sufficient to instigate a disruption. There is an initial cooling phase (pre-TQ) of 2.5ms, before a rapid (0.5ms) thermal quench (TQ), then a 24ms current quench. A summary of the experiment is shown in Fig. 1.

M3D-C¹ and simulation setup

Numerical simulations are performed using the extended-MHD code M3D-C¹ [1]. The initial condition is an EFIT reconstruction of the plasma equilibrium just before the disruption. Neon is added as a plume of 241 shards with a distribution of sizes and trajectories, and an average speed of 200m/s (see Fig. 2); its rate of ablation into the plasma is determined by a neutral gas shielding model [2] for each shard. Impurity ionisation, radiation and recombination is calculated for each charge state using the KPRAD model [3]. Spitzer resistivity is used, such that as the impurities cool the plasma edge, current is forced to diffuse inwards. We present results from two near-identical simulations — one in 2D, with material being deposited uniformly around an annulus, and one in 3D, with a Gaussian-shaped deposition of toroidal variance 2.4m. Our analysis focusses on the pre-TQ and identifying onset of the TQ due to MHD activity.



(a) Total radiated power from KB5V bolometer; simulation traces are shifted to start at 24.0315s, the timing of first light in the KL8-E8WA camera.



(b) Midplane electron temperature profile from KK3F ECE diagnostic.

Figure 3: Comparison of the pre-TQ phase between simulations and experiment.

Comparison of pre-TQ

During the pre-TQ, radiation increases with the quantity and temperature of ablated impurities as they penetrate. Measurement of total radiation from the KB5V bolometer, which is offset 90° toroidally from the SPI, is compared with simulations in Fig. 3a. The total is the right order of magnitude, despite only $\sim 4\%$ of the neon ablating during the simulations. The rapid growth indicative of TQ onset is well-captured in the 3D simulation (timing is relative to the observed first light at 24.0315s, not fitted to the data). This suggests the simulation may be capturing the correct MHD trigger, despite the fact it does not progress through the TQ. The two simulations agree well during the pre-TQ phase, with 3D effects responsible for the TQ radiation spike.

More detailed analysis of the pre-TQ phase is possible by looking at the electron temperature profile evolution from ECE measurements (Fig. 3b). Agreement with simulations is reasonable — the cooling front moves towards the core at the correct rate in both 2D and 3D cases, and the accelerated contraction as the TQ begins at 24.034s is reproduced in the 3D simulation.

Onset of MHD activity

In the 3D simulation, we observe progressive destabilisation of $2/1$ and $3/2$ MHD modes, with the latter heralding TQ onset (Fig. 4). Linearised stability analyses of snapshots from the 2D simulation can predict the timing of instability onset and yield consistent growth rates and mode shapes for $n = 1$ and 2 , suggesting axisymmetric current contraction is important in MHD destabilisation of the 3D case. However, a radiatively cooled $2/1$ island is also observed in 3D, the growth of which may contribute to TQ onset. The timing of $n = 1$ MHD activity is also consistent with experiment.




Figure 4: Onset of MHD activity in 3D simulation (top), 2D simulation (middle), and experiment (bottom). Inset are change in electron temperature in both simulations, change in flux function in 3D, and eigenmode in 2D.

Conclusions

Simulations of SPI into an Ohmic JET plasma using the extended-MHD code M3D-C¹, in both 2D and 3D, show a pre-TQ evolution of radiation and temperature profile consistent with experiment. The timing of MHD instability, indicated by an upturn in radiation and $n > 0$ magnetic energy, is also well-captured. Linear analyses of the 2D simulation predict instability compatible with the 3D results, although helical island cooling is also significant in 3D.

References

- [1] J. Breslau, N. Ferraro, S. Jardin, Some properties of the M3D-C¹ form of the three-dimensional magnetohydrodynamic equations. *Phys. Plasmas* **16**, 092503 (2009)
- [2] P. B. Parks, G. A. Gerdin, L. L. Vahala, A. G. Elcashlan, Model of ablation flow near light-atom pellets with surface boundary conditions. *Nucl. Fusion* **34**, 417–426 (1994)
- [3] B. C. Lyons, C. C. Kim, Y. Q. Liu, N. M. Ferraro, S. C. Jardin, J. McClenaghan, P. B. Parks, L. L. Lao, Axisymmetric benchmarks of impurity dynamics in extended-magnetohydrodynamic simulations. *Plasma Phys. Control. Fusion* **61**, 064001 (2019)

Acknowledgements

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014–2018 and 2019–2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission. Computing resources at PPPL and Marconi-Fusion (projects WPJET1, SPIM3D) are acknowledged. The assistance of Jin Chen and Seegyong Seol in installing M3D-C1 on Marconi, and Hongjuan Sun, Pedro Carvalho, Elena de la Luna, and Scott Silburn in JET data, is acknowledged. OPB is funded by a UKAEA-PPPL postdoctoral fellowship.