

UKAEA-CCFE-PR(20)88

A. R. Jackson, A. S. Jacobsen, K. G. McClements, C.
A. Michael, M. Cecconello

Diagnosing fast ion redistribution due to sawtooth instabilities using fast ion deuterium-alpha spectroscopy in the Mega Amp Spherical Tokamak

Enquiries about copyright and reproduction should in the first instance be addressed to the UKAEA Publications Officer, Culham Science Centre, Building K1/O/83 Abingdon, Oxfordshire, OX14 3DB, UK. The United Kingdom Atomic Energy Authority is the copyright holder.

The contents of this document and all other UKAEA Preprints, Reports and Conference Papers are available to view online free at scientific-publications.ukaea.uk/

Diagnosing fast ion redistribution due to sawtooth instabilities using fast ion deuterium-alpha spectroscopy in the Mega Amp Spherical Tokamak

A. R. Jackson, A. S. Jacobsen, K. G. McClements, C. A. Michael,
M. Cecconello

Diagnosing fast ion redistribution due to sawtooth instabilities using fast ion deuterium- α spectroscopy in the Mega Amp Spherical Tokamak

A. R. Jackson^{1,2}, A. S. Jacobsen², K. G. McClements², C. A. Michael³, M. Ceconello⁴

¹ Department of Physics, University of Durham, South Road, Durham, DH1 3LE, United Kingdom

² CCFE, Culham Science Centre, Abingdon, Oxfordshire, OX14 3DB, United Kingdom

³ Department of Physics and Astronomy, University of California - Los Angeles, Los Angeles, CA 90095-7099, USA

⁴ Uppsala University, P.O. Box 256, SE-751 05 Uppsala, Sweden

Abstract.

The efficient confinement of fast ions is crucial to the development of a fusion power plant. Sawtooth crashes are a manifestation of an internal kink instability, which can severely degrade this confinement. The effect of these on a fast ion population can be investigated using fast-ion deuterium- α (FIDA) spectroscopy, where charge exchange reactions between the fast ions and injected neutrals from a neutral beam injection (NBI) auxiliary heating system generate Doppler-shifted photons which can be observed and interpreted to provide information on the temporal evolution of the underlying fast-ion distribution function. Data obtained using this diagnostic can be interpreted using the codes TRANSP/NUBEAM and FIDASIM, which produces a synthetic spectrum for forward modelling of the experimental data. We have examined the effects of the sawtooth instability using these techniques on the Mega Amp Spherical Tokamak (MAST). Work reported by M. Ceconello *et. al.*, (Plasma Physics and Controlled Fusion 60, 055008 (2018)) showed that neutron camera data could not be used to distinguish between three models used in TRANSP to simulate the effects of sawteeth on the fast ions. We have extended this comparison between fast ion data and sawtooth models to FIDA measurements, and conclude that FIDA data also cannot be used to distinguish between the models. For FIDA channels that probe the sawtooth region, at each sawtooth crash there is an overall drop in the emission of up to 60%. Data from passive FIDA channels (i.e. with emission resulting from neutralisation by thermal neutrals in the plasma periphery rather than beam neutrals) show a sudden increase in the emission following sawtooth crashes. The subsequent decay in the emission in these passive channels indicate that redistributed passing fast ions are rapidly lost from the edge region, probably as a result of charge-exchange reactions with edge neutrals.

1. Introduction

Sawtooth crashes arise from a periodic MHD instability found in tokamaks, wherein an internal kink instability with dominant toroidal mode number $n = 1$ and dominant poloidal mode number $m = 1$ forms due to a peaking of the plasma current profile (equivalently, the safety factor q drops below 1 in the plasma core region). This instability is characterised by a sudden flattening of temperature and density profiles in the core of the plasma [1]. The sawtooth instability and its interaction with the fast ion population is of interest to the development of ITER plasmas, as crashes can degrade the quality of the fast ion confinement or more generally cause redistribution in both configuration and velocity space, in particular an expulsion of fast ions from the sawtooth region [2]. ITER plasmas rely on good fast ion confinement as the primary source of heating comes from the fusion α -particles. The presence of fast ions can increase the amplitude of sawtooth crashes by stabilising the instability, delaying the onset of the crash and therefore causing its consequences to be more violent when it does occur [3, 4]. Plasmas in the Mega-Amp Spherical Tokamak (MAST) were heated by deuterium neutral beams with primary particle energies of typically around 60–70 keV and thus contained substantial fast ion populations whose behaviour could be studied using a range of diagnostics. The MAST data investigated in this paper have been previously examined by a number of researchers [5, 6, 7]. The study presented here extends the work carried out by M. Cecconello *et. al.* [7] and utilises TRANSP/NUBEAM data generated in the course of that study. Specifically, we have extended the data analysis to include fast-ion deuterium- α (FIDA) data. The structure of this paper is as follows: in section 2 the background regarding the diagnostic and modelling software used in the analysis is given; in section 3 the models used to describe the effect of sawtooth crashes on the plasma are discussed; in subsection 4.1 the different models are compared using the experimental and synthetic FIDA data; in subsection 4.2 further analysis regarding the redistribution of the fast ions in the experiment is described and in section 5 some preliminary results regarding the analysis of passive FIDA emission are presented.

2. Fast-Ion Deuterium- α spectroscopy

The FIDA spectroscopic diagnostic can be utilised to gather local measurements of the fast-ion distribution present in the plasma. The neutral-beam injection (NBI) system introduces high energy deuterium neutrals into the plasma which can undergo charge exchange reactions with the fast-ion population (resulting from the ionisation of beam neutrals), generating a fast neutral that can be in an excited state. De-excitation from quantum state $n = 3$ to $n = 2$ generates a Balmer- α photon. The Doppler shift of this photon from the Balmer-alpha rest wavelength (656.1 nm) is proportional to the component of the fast-ion's velocity along the line of sight of the diagnostic [8]. The observed spectrum also has other *active* (present only when the neutral beam is on)

and *passive* (present whether the beam is on or off) spectral components that must be taken into account during interpretation. A component of the passive signal is FIDA light generated by charge exchange between cold edge neutrals and fast ions. Analysis of this is discussed further in section 5. This and other passive components such as bremsstrahlung can be subtracted from the raw signal. A toroidally displaced reference view (as was used on MAST [9]) observes plasma that does not contain a neutral beam. Assuming axisymmetry, this can be used to subtract from the main view to leave only the active signal. Alternatively, some tokamaks use beam notching, which requires instead the assumption of some degree of temporal stability. In order to interpret the data taken by the diagnostic, forward modelling is performed using the codes TRANSP/NUBEAM [10, 11] and FIDASIM [12]. TRANSP is a long-standing plasma transport and equilibrium solver that can be used to both analyse and predict plasmas. NUBEAM is a Monte-Carlo module for TRANSP that allows for the simulation of NBI systems and of the fast-ion populations that are generated. Profiles of a number of experimentally determined plasma parameters, such as (but not limited to) electron temperature and density, impurity density and toroidal plasma rotation at a particular time are given as input. A magnetic equilibrium is also required, which can be generated EFIT beforehand or can be evolved with TRANSP’s internal solver. The plasma is evolved in time and produces simulated fast-ion distributions, which form the crucial input to FIDASIM. FIDASIM uses these distributions to generate a synthetic active FIDA spectrum, which can then be compared with the background-subtracted experimental data. Figure 1 shows a synthetic spectrum generated with FIDASIM, indicating the form of a typical active signal as observed on MAST in terms of specific components.

Due to the specific geometry of the NBI system and the toroidally viewing FIDA diagnostic (the line-of-sight is co-current, as is the beam) the spectra is primarily red-shifted. The FIDA component is shown in red. The three peaks in yellow represents emission due to collisional excitation of injected neutrals (beam emission spectroscopy or BES). The location of the peak is dependent on the injection energy, with three peaks appearing due to there being a small proportion of di- and tri-atomic deuterium molecules in the NBI system, leading to full, half and third energy peaks. The halo component is generated by charge-exchange between the injected neutrals and the thermal ions. It may be noted that the halo component has a slight redshift, which is indicative of there being a bulk co-current rotation of the plasma. The MAST implementation of this diagnostic (described in detail in [9]) had both toroidal and vertical lines of sight, with associated passive views. Figure 2 shows the extent of the relevant toroidal views, along with the utilised NBI beamline and the last closed flux surface (LCFS). Due to the relative orientation of the lines-of-sight with respect to the magnetic field, toroidal views are primarily sensitive to passing particles, while vertical views are primarily sensitive to trapped particles.




Figure 1: Example spectra generated with FIDASIM of the toroidal view on MAST showing the FIDA, BES and halo components of the spectra. The dotted vertical line shows the location of the rest wavelength of deuterium- α emission, 656.1 nm. The BES (beam emission spectroscopy) peaks are caused by collisional excitation of beam neutrals to $n = 3$ states. There are three such peaks since the beam contains neutrals with energies equal to one half and one third of the primary energy.

3. Sawtooth models in TRANSP

The Kadomtsev model characterises the sawtooth crash as a process of rapid magnetic reconnection [13], in which equal and opposite helical flux on either side of the $q = 1$ surface (q being the safety factor) reconnect, forming a magnetic island that grows to become the new magnetic core of the plasma. The new core is colder and less dense than the original hot core, which is dispersed towards greater minor radii. In this model complete reconnection occurs - directly after the crash $q \geq 1$ everywhere in the plasma. Experimentally this model has been shown to be not applicable to all plasma pulses. Porcelli *et al.* [14] describe a number of pulses, some of which follow the Kadomtsev model and several that do not. The authors then propose an alternative model (referred hereafter as the Porcelli model) which allows for incomplete reconnection, meaning that $q(r = 0) < 1$. Here the formation of the magnetic island begins as in the Kadomtsev model but is interrupted by magnetic turbulence once the island has grown to some threshold size. The Kadomtsev model moves plasma at $q = 1$ pre-sawtooth to the magnetic axis after the sawtooth, and other regions are mixed with helical flux-matching as described above. This has a flattening effect on the temperature and density profiles, but can cause some previous structure to be retained. Alternatively, ergodic mixing can be utilised, ignoring the requirement of flux-matching and redistributing the plasma in such a way that the temperature and density profiles are




Figure 2: Plan view of MAST, showing an example last closed flux surface, along with the NBI beam and the relevant toroidal lines of sight.

completely flattened within the mixing region. The Porcelli model specifies two regions that are mixed separately, based on how large the magnetic island is allowed to grow before it is interrupted. Once the new locations for the markers have been determined, the post-crash equilibrium combined with conservation of magnetic moment and velocity parallel to the magnetic field determine the change in the energy and velocity of the markers [15]. TRANSP/NUBEAM implements the models in the KDSAW module, and a full explanation of the behaviour of the models is given in the online documentation [16]. Figure 3 shows synthetic spatial distributions of fast ions on MAST (generated by TRANSP/NUBEAM) before (Figure 3a) and after (Figure 3b) a sawtooth crash.

The three models were used in TRANSP/NUBEAM runs (provided by M. Cecconello *et. al.* [7]) to provide simulated fast-ion distributions before and after each sawtooth in pulse #29880. These were then utilised by the authors, with FIDASIM, to generate synthetic spectra, which are compared to the experimental data.

4. Analysis

The data used in this study derive from a series of three pulses (#29880-#29882) which were designed to be as nearly identical as possible. Figure 4 shows the time evolution of each pulse in a variety of parameters. The sawtooth crashes are clearly seen in the neutron rate and core electron temperature, but for these pulses the electron density is largely unaffected by the instability. To analyse the FIDA data, time slices were




Figure 3: The simulated fast-ion density in an azimuthal cross section of MAST before and after a sawtooth crash, with the flux surfaces shown in black.

taken before and after the third sawtooth in the overall sequence of sawteeth. The EFIT equilibrium reconstruction yields a q -profile greater than 1 at the time of the first sawtooth, and is not low enough after the second sawtooth to trigger the Porcelli model. A q substantially less than 1 is required to trigger the 3 sawtooth models in TRANSP/NUBEAM, a condition that is met by the time of the 3rd sawtooth in the sequence. Additionally, there is a secular downward trend in the experimental FIDA radiance over the course of the pulses, so the third sawtooth provides the best available signal-to-noise ratio. Figure 5 shows a core plasma soft x-ray camera trace for the selected single crash present in all three pulses. The soft X-ray emission can be regarded as a proxy for electron temperature.

The replication of the tokamak conditions between pulses is such that, in the sequence of crashes that occurs in all of the pulses, the selected crash is separated in time by a maximum of 7 ms between the pulses. This is significantly less than both the energy confinement time and the fast-ion slowing down time at around the time of the selected crash. From this it is concluded that the pulses are repeatable and that it is appropriate to average across pulses (after aligning the selected crashes in time) as well as averaging across time, thereby improving the signal-to-noise ratio. This technique was also used in the previous investigations [6, 7]. The BES peaks were extremely bright




Figure 4: Time traces of plasma parameters for the three pulses used in averaging. (a) NBI power, (b) neutron rate, (c) core electron temperature, (d) core electron density.




Figure 5: Soft X-ray (SXR) trace showing a sawtooth crash for shots #29880, #29881 and #29882.

compared to the FIDA light, which can be problematic as there can be pixel-to-pixel 'bleeding' in the spectrometer charge-coupled device (CCD) chip in such situations. An optical mask was used to block these features. Figure 6 shows an example of the full experimental spectrum along with the corresponding FIDASIM prediction (using the Kadomtsev model only, as an example).




Figure 6: Full experimental FIDA spectrum for pulse #29880 along with the corresponding synthetic spectrum, showing the summation of all the synthetic components and the synthetic FIDA component in isolation (using the Kadomtsev model only, as an example). The shaded area shows the extent of the experimental spectrum that is usable for analysis. All other wavelength regions are unusable for fast ion studies due to an optical mask blocking bright unwanted regions of the spectra, unwanted spectral components dominating the FIDA component, or the lack of any fast ion signal at all.

It can be seen that the presence of the other spectral components (and the mask) mean that there are no usable data in the lower end of the wavelength range, and the signal decays rapidly towards the upper end. What remains is the usable window, highlighted in green. The position and size of the window in wavelength space varies across different imaged positions in the plasma (spectrometer *channels*), but in every case there is a region of the spectrum dominated by the fast ions.

4.1. Comparison of different sawtooth models with FIDA data

In the work of Ceconello and co-workers [7] neutron camera data were insufficient to determine which of the three sawtooth crash models used (Kadomtsev, ergodic Kadomtsev and Porcelli) provided the best fit to the data. It was speculated that FIDA data may help in this regard. Figure 7 shows a before- and after-crash timeslice of the FIDA data, averaged over 3 ms (1–4 ms before crash, 1–4 ms after the crash) and across the three pulses (with the crash times aligned), along with synthetic spectra generated using each of the sawtooth models mentioned above, for a channel observing the plasma at a radius of 1.205 m in the plasma midplane. The synthetic data were only generated for pulse #29880. It is appropriate to do this for the reasons given at the beginning of this section, i.e. that the plasma conditions are closely replicated in the three pulses that are analysed 4.




Figure 7: Experimental and synthetic spectra showing only the region where FIDA light can be observed at $R=1.025$ m in the plasma midplane. The predictions based on the three models are shown.

Before the crash there is a difference between the predicted spectra. As mentioned previously, due to the equilibrium reconstruction used, none of the models trigger at the time of the first sawtooth, and only the Kadomtsev and Ergodic Kadomtsev model trigger at the second sawtooth. Therefore we observe a small difference in the models before the crash. Drops in spectral radiance are observed in both the experimental and synthetic data for the channel shown in Figure 7 and for channels closer to the core, with channels further out showing small to no drop. The absolute values for the spectral radiance is reproduced well by the TRANSP/NUBEAM and FIDASIM modelling, for the shown channel in Figure 7. However, this degree of agreement is only

seen in the shown channel; other channels show serious deviation between experiment and prediction. This discrepancy in the absolute magnitude, possibly due to calibration factors, is discussed further in section 4.2. In all channels, over most of the usable wavelength space, the 3 sawtooth models produce only a small difference in the predicted spectra compared to the overall change due to the sawtooth. The errors shown are based on the combination of photon shot noise and the CCD read noise for both the active and passive channels used (dark current noise is neglected due to the cooling of the CCD chip). It must be emphasised that this represents the minimum possible error on the FIDA data and that the true error may be larger, especially considering the possible problems due to uncertain calibration, discussed in subsection 4.2. Due to this uncertainty and the size of the error, no clear determination of the suitability of one model over the others can be made. We conclude that FIDA measurements cannot resolve the issue of which model best represents the data [7].

4.2. Detailed analysis of fast-ion redistribution by sawtooth crash and comparison with FIDA

While in the previous section it was shown that FIDA is not sensitive enough to distinguish between the 3 sawtooth models, and that there was significant deviation of the experiment and prediction, by eliminating a source of calibration error it is shown in this section that we can achieve good agreement between the forward-modelled FI redistribution due to the sawtooth crash and the diagnostic within the experimental error. Only some of the FIDA channels contain usable spectral data. Channels close to the plasma edge view plasma regions containing relatively few fast ions and those close to the plasma core view plasma regions containing few neutrals with which the fast ions can charge exchange. Both of these effects lead to a low level of FIDA radiance. Of the channels with a usable level of FIDA radiance it was observed that only a single channel, observing the plasma at a major radius of 1.205 m, consistently showed agreement between the absolute experimental values and the FIDASIM predictions. However, it has been shown previously that a scaling factor is required in order to model the experimental BES peaks correctly [9]. Applying that factor here would lead to larger disagreement between the model and data, and would have varying effects on the agreement in the other channels. Due to the presence of the optical mask it is not possible to use the comparison of the experimental and synthetic BES peaks here as a test of the calibration and scaling validity. Investigating relative changes in the spectra between pre- and post-crash reduces or eliminates some of the effects of absolute miscalibration. Figure 8 shows the relative change (difference in signal after the crash vs before, divided by the signal before) in the observed spectral radiance across the fast-ion dominated portion of the spectrum, across channels that have an observable level of FIDA radiance averaged using the same methods described for Figure 7 above.

The location of the BES peaks in wavelength space are dependent on the relative orientations of the line-of-sight to the NBI beamline, so they appear at different locations




Figure 8: Spectral radiance across a number of FIDA channels. Note that the full energy BES peak appears at different locations depending on channel. The grey shaded area in plots (a)-(e) designates where the effect of the peak has a significant effect on the synthetic data, so direct comparisons between the observations and the models cannot be made here.

in wavelength for the different FIDA channels, which can reduce the usable areas of the spectrum. Unusable parts of the spectra due to this effect are shown by the greyed out portions in figure 8. These were determined by calculating the relative change in the synthetic spectra before and after the sawtooth crash firstly using all FIDASIM spectral components and again with only the FIDA component. The unusable region was set to where the difference of these calculations was greater than 0.5% (in reference to the y-axis scale of the plot). Note as well that the error on the experimental data increases significantly at wavelengths above 661.0 nm, as the absolute signal experiences a rapid drop-off. It can be seen that in the valid region the experimental and synthetic results are comparable for the most part except at 1.139 m (Figure 8(d)), where there is significant disparity between them. Through examination of Thomson scattering data, it can be determined that the outboard radius of the inversion region in the midplane is approximately 1.16 m, slightly higher than the tangency radius corresponding to Figure 8(d) and slightly lower than that corresponding to Figure 8(e). Comparing these two plots, it can be seen that the relative change in emission following the sawtooth crash is considerably larger in Figure 8(d) than it is in Figure 8(e). This in itself is unsurprising, since the effects of the sawtooth on the emission are expected to become smaller as the line of sight moves away from the sawtooth region. The fact that an effect is seen at all in Figure 8(e), despite the tangency radius lying outside the sawtooth region, can be explained by the fact that the fast ion orbit widths and Larmor radii are typically larger than the difference between the tangency radius of this spectrometer chord (1.205 m) and

the sawtooth inversion radius (1.16 m). When we compare the drops in FIDA emission with the three sawtooth models, it is evident from Figure 8(d) that all of them predict a drop in emission that is smaller than the measured value. In making these comparisons, it should be noted that while finite Larmor radius corrections are taken into account in NUBEAM for the purpose of evaluating neutralisation probabilities and collision rates, they are not taken into account in terms of the local magnetic field used to track particle orbits, since a guiding centre approximation is used [11]. This introduces an error in the synthetic spectra for tangency radii lying close to the boundary of the sawtooth region. The discrepancy between measurements and modelling in Figure 8(d) could perhaps be due to a pitch angle (i.e. Larmor radius) dependence of the effects of the sawtooth on the fast ion population that is not captured by any of the models, although further investigation would be needed to confirm this. Figure 8(d) shows the largest drop in the observed FIDA radiance across all valid channels, along with the largest drop in the Ergodic Kadomtsev and Porcelli models. Figure 8(e) shows the largest drop in the Kadomtsev model, with similar drops in the other models to Figure 8(d). This is somewhat counter-intuitive: it may be thought that channels corresponding to locations closer to the core would have larger drops in radiance, since the core is inside the sawtooth region and contains more fast ions prior to the crash. The drop in radiance in the channel used in Figure 8(e) and the lesser drops in other channels can be correlated with the change in the fast-ion distributions predicted by TRANSP/NUBEAM due to the good correspondence between the observed and predicted values. Figure 9 shows the computed fast-ion distributions at a major radius of 1.205 m in the midplane of the tokamak (the Kadomtsev model is used here, as an example). Pitch here is defined as the ratio of the component of the marker's velocity parallel to the magnetic field to the full velocity. On MAST during conventional operation the toroidal component of the magnetic field, B_ϕ , was directed clockwise as seen from above and current anti-clockwise, so particles with negative pitch are travelling in the co-current direction. The overlaid contours show the extent of the *weight function* for the FIDA diagnostic [9, 17], specifically in the chosen spatial location and over the range of wavelengths that are dominated by the presence of the fast ions. Weight functions are essentially a measure of the probability that a CX reaction emits a photon of a specific wavelength (or range of wavelengths), given the velocity of the fast ion and the angle of the diagnostic line-of-sight to the local magnetic field [18, 19].

For comparison with Figure 8(e) the relative change in the fast-ion distributions across the three models is shown in Figure 10.

It can be observed across all of the models in Figure 10 that there is a significant loss of fast ions (up to 50%) from energies between 45–55 keV across pitches between -0.25 and -1 , with the Kadomtsev (Figure 10(a)) model showing additional losses down to 30 keV and the Ergodic Kadomtsev (Figure 10(b)) and Porcelli (Figure 10(c)) models showing an increase in the fast ions around a pitch of -0.75 across energies 25–40 keV. Additionally there is an increase in fast ions with energies less than 20 keV across positive pitch values. The higher energy particles are expelled from the sawtooth region




Figure 9: Simulated fast ion distributions for $R = 1.205\text{m}$ in the device midplane using the Kadomtsev model. (a) Before crash, (b) after crash. The green contours represent the weight function for the FIDA diagnostic in the fast-ion dominated wavelength range and the spatial location of measurement.

during the crash and lower energy particles from larger radii are mixed in. These lower energy fast ions will have undergone significant pitch angle scattering. The apparent preference for positive pitch angle after the crash across the models in Figure 10 is likely to be an artefact of the normalisation. It can be seen that the weight function primarily intersects the region of significant loss, which is reflected in the decrease in FIDA radiance shown in Figure 8, with the smaller regions of increase within the intersection present in the Ergodic Kadomtsev and Porcelli models leading to the smaller decrease present in those models compared to the Kadomtsev. The shape of the weight function depends on the tangency radius of the line of sight. In the other channels the weight function primarily intersects the population of newly injected fast ions around 55 keV, so in these channels a smaller change in FIDA radiance due to redistribution of the FI population by the sawtooth is observed. It should be emphasised when examining synthetic fast-ion distributions that any regions of the phase-space outside of the overlap with the weight function cannot be probed with the FIDA diagnostic, so cannot be compared with experimental values. The recovery of the fast-ion distribution after the crash can be observed in Figure 11, where the spectra move towards pre-crash conditions as the time window averaged for the 'after' portion of the data moves further from the




Figure 10: The relative change in the fast-ion distribution function for $R = 1.205\text{m}$ in the device midplane. (a) Kadomtsev, (b) Ergodic Kadomtsev, (c) Porcelli. Areas in green represent phase-space regions with a fast-ion presence after the crash that had no presence before. Note that in some regions there is a positive change of greater than 100%. The black contours represent the weight function for the FIDA diagnostic in the fast-ion dominated wavelength range and the spatial location of measurement.

crash. The total time interval covered in this figure is comparable to the fast-ion slowing down time, which is the timescale on which we could expect the fast-ion distribution to recover following a sawtooth crash. Figure 4(b) shows an overall degradation of the fast-ion confinement, which could contribute to the incomplete recovery of the spectra. There is a significant disagreement between the experimental and synthetic data in Figure 11(e). While we do not yet have a definitive explanation for this, examination of the $D\text{-}\alpha$ signal for one of the three pulses (#29880) in the period after the third sawtooth shows a significant increase around the start of this averaging window. This could be a consequence of more edge neutrals permeating into the hot plasma, which would be likely to generate an increase in the passive FIDA signal, thereby affecting the passive subtraction. Such an effect was not taken into account in the FIDASIM calculation for these pulses.

5. Passive FIDA

As described in section 2 there is a FIDA spectral component generated by charge exchange reactions between fast ions and neutrals present in the colder edge region of the plasma, rather than the beam neutrals. In this section, the effect of the sawtooth crashes (in #29880) on the passive emission observed by a toroidally directed passive view on MAST is examined. Figure 12a shows a plan view of the MAST plasma, with the black circles denoting the minimum and maximum extent of the plasma at the




Figure 11: Relative change in the spectral radiance at $R = 1.205\text{m}$ in the device midplane, averaged over the indicated time windows. Note that Figure 11(a) examines the same data as Figure 8(e)

chosen time. The fan of chords available to the passive FIDA view are shown in grey, with an example channel shown in red. Figure 12b shows the integrated radiance, I , for this channel for a single sawtooth crash. A decaying exponential of the form:

$$I(t) = ae^{-\frac{(t-t_0)}{\tau}} + c \quad (1)$$

where t is time, t_0 is a time offset, and a , c and τ are constants, is used to describe the radiance temporal decay. We fit the decay time τ to the decaying part of the signal. It is observed that there is a large spike in integrated radiance across all channels that intersect the plasma, with decay times between 0.3–0.5 ms. This feature is not observed in data integrated in regions free from FIDA light, indicating that background fluctuations are not the cause. Therefore the data suggests that there is a sudden significant increase in the population of passing fast ions followed by a rapid decrease. It is likely that this rapid decrease is due primarily to charge exchange between these passing fast ions and edge neutrals rather than direct fast ion orbit losses. A typical neutral density, n_0 , in the MAST plasma edge is a few times $1 \times 10^{16} \text{ m}^{-3}$ [20]: using the velocity, v , and charge exchange cross-section σ for a 40 keV deuteron [21], we estimate the charge-exchange time to be $\tau_{CX} = 1/(vn_0\sigma) \sim 0.3 \text{ ms}$, consistent with the decay time in Figure 12b. Bolte *et. al.* [22] have carried out a detailed examination of passive FIDA emission on the DIII-D tokamak, where they also observe the preference for the ejection of passing particles from the core to the edge. Additionally, they are able to determine, with an accurately calibrated diagnostic and the use of passive FIDASIM simulations, that approximately 1% of the fast-ion population is expelled into the edge region. Looking to the future, MAST-Upgrade will have a fast-ion loss detector, which




Figure 12: (a) Available toroidally-viewing passive chords, with selected chord highlighted in red, and the NBI beamline to emphasise the passive nature of the view. Data from this chord is shown in Figure 12b. The black circles show the minimum and maximum extent of the plasma. (b) Spike and decay of passive signal seen during a sawtooth crash. The decaying part of the signal has been fitted with an exponential.

is expected to provide direct information on sawtooth-induced fast-ion losses.

6. Conclusion

Based on forward modelling performed with the FIDASIM code in conjunction with experimental FIDA data it is not possible to distinguish between complete (Kadomtsev and ergodically mixed Kadomtsev) and incomplete (Porcelli) reconnection models for the sawtooth crashes, due to the level of noise present in the FIDA data and the relatively small differences seen in the FIDASIM data between the models. While issues regarding the calibration of the diagnostic mean that there are significant difficulties in analysing the absolute FIDA radiances, examining the relative change in the spectra and synthetic fast-ion distribution resulting from the sawtooth crashes allows for further investigation into the fast-ion redistribution. Large drops in the measured spectral radiance are reflected in the synthetic spectra, which indicates significant depletion of the fast-ion phase-space at energies around 40 keV, which progressively recovers over the inter-sawtooth period. Passive data show that sawtooth crashes lead to a sudden increase and subsequent rapid decay of passing fast ions in the edge region of the plasma. We conclude that additional fast ion diagnostics, such as those that will be available in MAST-Upgrade, are likely to be required in order to discriminate experimentally between alternative models of fast ion redistribution and loss resulting from sawtooth

crashes.

Acknowledgements

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission. Funding from EP-SRC grant number EP/T012250/1 is acknowledged.

- [1] S. von Goeler, W. Stodiek, and N. Sauthoff, “Studies of Internal Disruptions and $m=1$ Oscillations in Tokamak Discharges with Soft—X-Ray Techniques,” *Physical Review Letters*, vol. 33, no. 20, pp. 1201–1203, 1974. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.33.1201>
- [2] F. Wising, D. Anderson, and M. Lisak, “Modelling of fast ion redistribution due to sawteeth in neutral beam heated plasmas,” *Plasma Physics and Controlled Fusion*, vol. 34, no. 5, pp. 853–862, 1992.
- [3] I. T. Chapman, S. D. Pinches, J. P. Graves, R. J. Akers, L. C. Appel, R. V. Budny, S. Coda, N. J. Conway, M. De Bock, L. G. Eriksson, R. J. Hastie, T. C. Hender, G. T. Huysmans, T. Johnson, H. R. Koslowski, A. Krämer-Flecken, M. Lennholm, Y. Liang, S. Saarelma, S. E. Sharapov, and I. Voitsekhovitch, “The physics of sawtooth stabilization,” *Plasma Physics and Controlled Fusion*, vol. 49, no. 12 B, pp. B385–B394, 2007.
- [4] I. T. Chapman, R. J. Buttery, S. Coda, S. Gerhardt, J. P. Graves, D. F. Howell, A. Isayama, R. J. La Haye, Y. Liu, P. Maget, M. Maraschek, S. Sabbagh, and O. Sauter, “Empirical scaling of sawtooth period for onset of neoclassical tearing modes,” *Nuclear Fusion*, vol. 50, no. 10, p. 102001, 2010.
- [5] B. Madsen, M. Salewski, J. Huang, A. S. Jacobsen, O. Jones, and K. G. McClements, “Velocity-space tomography using prior information at MAST,” *Review of Scientific Instruments*, vol. 89, no. 10, pp. 1–6, 2018. [Online]. Available: <http://dx.doi.org/10.1063/1.5035498>
- [6] M. Cecconello, O. M. Jones, W. U. Boeglin, R. V. Perez, D. S. Darrow, I. Klimek, S. E. Sharapov, M. Fitzgerald, K. G. McClements, D. L. Keeling, S. Y. Allan, C. A. Michael, R. J. Akers, N. J. Conway, R. Scannell, M. Turnyanskiy, and G. Ericsson, “Energetic ion behaviour in MAST,” *Plasma Physics and Controlled Fusion*, vol. 57, no. 1, p. 014006, 2015.
- [7] M. Cecconello, A. Sperduti, and the MAST team, “Study of the effect of sawteeth on fast ions and neutron emission in MAST using a neutron camera,” *Plasma Physics and Controlled Fusion*, vol. 60, no. 5, p. 055008, 2018.
- [8] W. W. Heidbrink, “Fast-ion $D\alpha$ measurements of the fast-ion distribution,” *Review of Scientific Instruments*, vol. 81, no. 10, p. 10D727, 2010.
- [9] C. A. Michael, N. Conway, B. Crowley, O. Jones, W. W. Heidbrink, S. Pinches, E. Braeken, R. Akers, C. Challis, M. Turnyanskiy, A. Patel, D. Muir, R. Gaffka, and S. Bailey, “Dual view FIDA measurements on MAST,” *Plasma Physics and Controlled Fusion*, vol. 55, no. 9, p. 095007, 2013.
- [10] R. J. Hawryluk, “An empirical approach to tokamak transport,” in *Physics of Plasmas Close to Thermonuclear Conditions*, 1981, pp. 19–46.
- [11] A. Pankin, D. McCune, R. Andre, G. Bateman, and A. Kritz, “The tokamak Monte Carlo fast ion module NUBEAM in the national transport code collaboration library,” *Computer Physics Communications*, vol. 159, no. 3, pp. 157–184, 2004.
- [12] W. W. Heidbrink, D. Liu, Y. Luo, E. Ruskov, and B. Geiger, “A code that simulates fast-ion $D\alpha$

- and neutral particle measurements,” *Communications in Computational Physics*, vol. 10, no. 3, pp. 716–741, 2011.
- [13] B. B. Kadomtsev, “On the tearing instability in tokamaks,” *Fizika Plazmy*, vol. 1, no. 5, pp. 710–714, 1975.
- [14] F. Porcelli, D. Boucher, and M. Rosenbluth, “Model for the sawtooth period and amplitude,” *Plasma Physics and Controlled Fusion*, vol. 38, pp. 2163–2186, 1996.
- [15] M. Podesta, “Private communication,” 2019.
- [16] D. McCune, “NTCC module – kdsaw – Kadomtsev Sawtooth Model with Porcelli Upgrade.” 2005. [Online]. Available: <https://w3.pppl.gov/ntcc/KDSAW/DOC>
- [17] W. W. Heidbrink, Y. Luo, K. H. Burrell, R. W. Harvey, R. I. Pinsker, and E. Ruskov, “Measurements of fast-ion acceleration at cyclotron harmonics using Balmer-alpha spectroscopy,” *Plasma Physics and Controlled Fusion*, vol. 49, no. 9, pp. 1457–1475, 2007.
- [18] M. Salewski, B. Geiger, D. Moseev, W. W. Heidbrink, A. S. Jacobsen, S. B. Korsholm, F. Leipold, J. Madsen, S. K. Nielsen, J. Rasmussen, M. Stejner, and M. Weiland, “On velocity-space sensitivity of fast-ion D-alpha spectroscopy,” *Plasma Physics and Controlled Fusion*, vol. 56, no. 10, 2014.
- [19] A. S. Jacobsen, “Methods to determine fast-ion distribution functions from multi-diagnostic measurements,” Ph.D. dissertation, Technical University of Denmark, 2015.
- [20] K. G. McClements, K. Tani, R. J. Akers, Y. Q. Liu, K. Shinohara, H. Tsutsui, and S. Tsujii, “The effects of resonant magnetic perturbations and charge-exchange reactions on fast ion confinement and neutron emission in the Mega Amp Spherical Tokamak,” *Plasma Physics and Controlled Fusion*, vol. 60, no. 9, 2018.
- [21] J. Wesson, *Tokamaks*, 3rd ed. Oxford: Oxford University Press, 2004.
- [22] N. G. Bolte, W. W. Heidbrink, D. Pace, M. Van Zeeland, and X. Chen, “Measurement and simulation of passive fast-ion D-alpha emission from the DIII-D tokamak,” *Nuclear Fusion*, vol. 56, no. 11, 2016.